

## Metabolic Labeling of Yeast RNA with Radioactive Uracil

Yuehua Wei\*

No.3 People's Hospital, Shanghai Jiao Tong University, Shanghai, China

\*For correspondence: [weiyh.sjtu.edu@gmail.com](mailto:weiyh.sjtu.edu@gmail.com)

**[Abstract]** To examine gene expression, northern blot or Real-Time PCR can be used to detect low abundant RNA such as mRNA. However, for high abundant RNAs such as rRNA and tRNA, Northern blot will not be able to discriminate the newly synthesized RNA from total RNA. Therefore, metabolic labeling is necessary to evaluate the expression of rRNA and tRNA genes. In this protocol, I describe a step-by-step method for labeling yeast RNA with radioactive uracil and examine the synthesis of these high abundant RNAs.

### Materials and Reagents

1. Yeast strain of interest
2. RapidGel (500 ml) (Affymetrix, catalog number: 75848)
3. Urea ( $\text{CO}(\text{NH}_2)_2$ ) (Sigma-Aldrich, catalog number: U6504)
4. Dextrose ( $\text{C}_6\text{H}_{12}\text{O}_6$ ) (Sigma-Aldrich, catalog number: G7021)
5. Yeast nitrogen base without amino acids (Sigma-Aldrich, catalog number: Y0626)
6. Synthetic dropout supplement without uracil (Sigma-Aldrich, catalog number: Y1501)
7. Uracil ( $\text{C}_4\text{H}_4\text{N}_2\text{O}_2$ ) (Sigma-Aldrich, catalog number: U1128)
8. Rapamycin
9. TEMED/Tetramethylethylenediamine ( $\text{C}_6\text{H}_{16}\text{N}_2$ ) (Thermo Fisher Scientific, catalog number: 110-18-9)
10. Ammonium persulfate (APS) ( $(\text{NH}_4)_2\text{S}_2\text{O}_8$ ) (Sigma-Aldrich, catalog number: A3678)
11. Formamide ( $\text{CH}_3\text{NO}$ ) (Thermo Fisher Scientific, catalog number: 75-12-7)
12. DEPC/ Diethylpyrocarbonate ( $\text{O}(\text{COOC}_2\text{H}_5)_2$ ) (Sigma-Aldrich, catalog number: D5758)
13. Bromophenol Blue ( $\text{C}_{19}\text{H}_{10}\text{Br}_4\text{O}_5\text{S}$ ) (Sigma-Aldrich, catalog number: B0126)
14. Xylene Cyanol FF ( $\text{C}_{25}\text{H}_{27}\text{N}_2\text{NaO}_6\text{S}_2$ ) (Sigma-Aldrich, catalog number: X4126)
15. [5, 6-<sup>3</sup>H]-Uracil (PerkinElmer, catalog number: NET368250UC)
16. RNA marker (Promega Corporation, catalog number: G3191)
17. Tris base (Thermo Fisher Scientific, catalog number: 77-86-1)
18. Boric acid (Thermo Fisher Scientific, catalog number: 10043-35-3)
19. EDTA (Sigma-Aldrich, catalog number: EDS-1KG)
20. Formamide loading dye

21. Small RNA separating gel
22. SD-Ura- (see Recipes)
23. SD-1/3 Uracil (see Recipes)
24. DEPC water (see Recipes)
25. 10x TBE in DEPC water (see Recipes)
26. Small RNA separating gel (see Recipes)
27. 2x Formamide loading dye (see Recipes)

### **Equipment**

1. Bench top centrifuge
2. 30 °C Shaker
3. Power supply with constant voltage > 450 V
4. Gel dryer (Bio-Rad Laboratories, catalog number: 165-1745)
5. Exposure cosset/intensifier screen (Sigma-Aldrich, catalog number: C5479-1EA)
6. 50 ml conical tubes
7. 1.5 ml Eppendorf tube

### **Procedure**

1. Inoculate yeast single colony in SD medium (or SD with appropriate dropouts). Shake 300 rpm at 30 °C overnight.
2. Dilute overnight culture in 50 ml conical tubes to 10 ml, OD<sub>600</sub> =0.1 with SD-1/3 Uracil and continue shaking until OD<sub>600</sub> =0.4 (*note: Reduction in cold uracil will allow hot uracil to be taken up by cells easily*).
3. Cells were treated with drug and control vehicle, for example 100 nM rapamycin (final concentration) and its solvent methanol, continue shaking 300 rpm at 30 °C for desired time. In this specific experiment, rapamycin were added for 30 min.
4. Collect yeast cells by spinning down at 1,000 x g for 1 min at room temperature, remove supernatant (*critical: Avoid putting yeast cells on ice. This is because ice will slow down growth, which will reduce significantly the uptake of hot uracil in the step 8 below*).
5. Re-suspend yeast cells in 1 ml SD-Ura- (*critical: Pre-warm medium to 30 °C*), transfer to 1.5 ml Eppendorf tube.
6. Spin down briefly by a bench top centrifuge at 5,000 x g, 15 sec, remove supernatant. Re-suspend yeast cells in 1 ml pre-warmed SD-Ura-.

7. Repeat 6 for 2 times and with the final re-suspension in 0.5 ml pre-warmed SD-Ura- (from the next step, collect radioactive liquid and solid waste in all steps, dispose according to environmental regulation).
8. Carefully add [5, 6-<sup>3</sup>H]-Uracil into each tube to the final concentration of 15 µCi/ml, vortex to mix, then put on a rack at 30 °C for 5 min.
9. Briefly spin down by a bench top centrifuge at 5,000 x g, 15 sec, remove supernatant.
10. Wash cells with SD-Ura-3 times as in step 5, ready to extract total RNA.
11. Total RNA was extracted by hot phenol method described in Wei (2012).
12. RNAs can be stored at -80 °C for up to 6 months.
13. Prepare “small RNA separating gel” on a large gel set (around 20 x 30 cm, mini gel did not work well).
14. Pre-run the gel for about 1 h at constant 450 V until the gel is heated to 50 °C.  
*Note: I found this step to be critical. One reason could be that pre-running the gel to this temperature could help get rid of excessive urea in the gel, making RNA possible to go through. I usually attached a thermometer to ensure that the temperature has reached 50 °C.*
15. Mix RNA samples with “2x Formamide loading dye” and heat at 70 °C for 2 min, then put samples on ice.  
Turn off power. Rinse the wells with 1x TBE using a syringe with needle, make sure all urea is rinsed out from the wells.  
*Note: Urea is very dense and it will be impossible to load samples if urea is not rinsed out. If residual urea remains in the well, the resulting bands will be wavy.*
16. Load RNA samples (25 µg) with appropriate RNA marker and run gel at constant 450 V for about 2 h (BPB runs around 12 nt and cyanol around 55 nt).
17. Stain the gel with EtBr and take picture under UV light. This is total RNA which served as controls for newly synthesized RNA.
18. Sandwich the gel with autoclaved filter paper on one side and Saran-Wrap on the other side, put on a gel-dryer with filter paper side attach to the vacuum surface. Dry the gel at 80 °C for at least 2 h. Sometimes gels get cracked, may be because of insufficient drying or leaky vacuum.
19. The dried gel will stick to the filter paper. Wrap them in Sara-Wrap. <sup>3</sup>H autoradiography in an exposure cassette with appropriate intensifying screen, for example Sigma Transcreen.
20. Develop after 4-7 days of exposure.

## Recipes

1. SD-Ura-  
 Synthetic dextrose medium with Uracil dropout:  
 20 g Dextrose  
 1.7 g Yeast Nitrogen Base  
 1.92 g synthetic dropout supplement without Uracil  
 5.0 g Ammonium Sulfate  
 Add ddH<sub>2</sub>O<sub>2</sub> to 1 L and autoclave.
2. SD-1/3 Uracil  
 SD medium with 1/3 of regular Uracil  
 Similar to SD-Ura- except adding 25 mg Uracil.
3. DEPC water  
 Add 1 ml DEPC to 1 L ddH<sub>2</sub>O<sub>2</sub>, mix and put at room temperature overnight. Autoclave.
4. 10x TBE in DEPC water  
 In 800 ml DEPC water, add:  
 108 g Tris base  
 55 g boric acid  
 40 ml of 0.5 M EDTA (pH 8.0)  
 Mix to dissolve and add DEPC water to 1 L. Autoclave.
5. Small RNA separating gel
  - a. Mix 2.5 ml 10x TBE, 6.25 ml RapidGel (40%) and 15 g Urea, heat to 50 °C and mix to dissolve.
  - b. Add DEPC water to 25 ml then filter through 0.45 µM Syringe.
  - c. Add 25 µl TEMED and 50 µl 25% APS, mix vigorously transfer to gel set with appropriate comb.
6. 2x Formamide loading dye  
 95% (v/v) formamide in DEPC water, add tiny amount of Bromophenol Blue (0.01~0.1%) and Xylene Cyanol FF (0.01~0.1%), vortex to mix.

## Acknowledgments

This protocol is derived from the following papers, Wei *et al.* (2009a) and Wei *et al.* (2009b), and the relevant references therein. The work was supported by NIH grants R01-CA099004 and R01-CA123391 to Dr. X.F. Steven Zheng at Robert Wood Johnson Medical School, University of Medicine and Dentistry of New Jersey.

**References**

1. Wei, Y., Tsang, C. K. and Zheng, X. F. (2009a). [Mechanisms of regulation of RNA polymerase III-dependent transcription by TORC1](#). *EMBO J* 28 (15): 2220-2230.
2. Wei, Y. and Zheng, X. F. (2009b). [Sch9 partially mediates TORC1 signaling to control ribosomal RNA synthesis](#). *Cell Cycle* 8(24): 4085-4090.
3. Wei, Y. (2012). [A Simple Preparation of RNA from Yeast by Hot Phenol for Northern Blot](#). *Bio-protocol* 2(9): e209.