

Biochemical Assays for MTase Activity

Yu Chen* and Deyin Guo*

College of Life Sciences, Wuhan University, Wuhan, China

*For correspondence: chenyu@whu.edu.cn; dguo@whu.edu.cn

[Abstract] Methyltransferase (MTase) transfers a methyl group (-CH₃) from the donor S-adenosyl-L-methionine (AdoMet or SAM) to biologically active molecules such as hormones, neurotransmitters, lipids, proteins and nucleic acids. The addition of a methyl group causes a change in the physicochemical properties of the molecules. The mRNA cap structure is essential for cell and virus. Guanine-N7-methyltransferase (N7-MTase) methylates the GpppN cap at the N7 position of guanine, resulting in cap-0 structure (m7GpppN), and Ribose 2'-O-MTase further methylates the first nucleotide of higher eukaryotic cellular and viral mRNAs at the ribose 2'-OH position to form cap-1 (m7GpppNm) structures. Here, we describe a biochemical assay to detect the activities of mRNA capping MTases.

Materials and Reagents

1. Bodicon m7G capping system (Bodicon, catalog number: CS0130)
2. S-adenosyl methionine (SAM) (involved in Bodicon m7G capping system) (Bodicon, catalog number: CS0130)
3. Bodicon Capping Enzyme (10 U/μl) (involved in Bodicon m7G capping system) (Bodicon, catalog number: CS0130)

Note: Because the sale of this kit was low, the previous companies which provide this capping kit were out of service. This capping kit was provided by a new company in China as custom-made products (contact e-mail: service@bodicon.cn, phone: +86-13628662011). In fact the similar capping kit from any other companies (such as Epicentre Biotechnologies, ScriptCap m7G capping system, catalog number: SCCE0610) is suitable for this experiment, and people can also contact with us to get the related protein or kit.

4. Inorganic pyrophosphatase (YIPP) (New England Biolabs, catalog number: M2403S)
5. S-adenosyl [methyl-³H] methionine ([³H]-SAM) (PerkinElmer, catalog number: NET155H001MC)
6. DEAE Sephadex (GE Healthcare, catalog number: 17-0170-01)
7. GTP (Thomas Scientific, catalog number: R0461)
8. RNase inhibitor (Thomas Scientific, catalog number: EO0381)
9. RNase free water
10. Phenol-chloroform (pH 4.8-5.2 for RNA only)
11. Ethanol (RNase free)

12. RNase free water
13. Sodium Dodecyl Sulfonate (SDS)
14. Ethylene Diamine Tetraacetic Acid (EDTA)
15. NH_4HCO_3
16. NaCl
17. 10x MTase Buffer (see Recipes)
18. Cap-0 cap structure (m7GpppN-RNA) (see Recipes)
19. Non-methylated Cap-0 cap structure (GpppN-RNA) (see Recipes)
20. MTase assay reaction mix (see Recipes)

Equipment

1. Bechtop
2. Water bath
3. Centrifuge
4. Liquid scintillation counter

Procedure

1. Synthesis of RNA cap structure as substrates.

The nascent RNA transcribed *in vitro* possesses a 5' triphosphate end. Two RNA capping systems are used to synthesize cap structure as the substrates of MTase.

- a. Combine 1-10 μg RNA and RNase free water up to 12.5 μl of total reaction volume.
- b. To heat to denature the *in vitro* transcribed RNA, incubate the tube at 65 °C for 10 min, and transfer the tube to ice immediately.
- c. Combine the following reaction components in the order given:

Cap-0 cap structure (m7GpppN-RNA)

Heat denatured RNA	12.5 μl
10x Bodicon Capping Buffer	2 μl
10 mM GTP	2 μl
3 mM SAM	1 μl
RNase inhibitor 40 U/ μl	0.5 μl
Bodicon Capping Enzyme (10 U/ μl)	2 μl
Total	20 μl

Non-methylated Cap-0 cap structure (GpppN-RNA)

Heat denatured RNA	12.5 μl
10x Bodicon Capping Buffer	2 μl
10 mM GTP	2 μl

Inorganic pyrophosphatase 0.1 U/μl	1 μl
RNase inhibitor 40 U/μl	0.5 μl
Bodicon Capping Enzyme (10 U/μl)	2 μl
Total	20 μl

- d. Incubate at 37 °C for 2 h.
- e. Purify the RNA substrates by using phenol-chloroform extraction and ethanol precipitation methods.
2. Prepare 10x MTase Buffer. The MTase buffer may be changed depending on different interested MTases.
3. Combine the following reaction components in the order given for MTase assays:

Purified MTases	1 μg
10x MTase Buffer	3 μl
GpppN-RNA or m7GpppN-RNA	2 μg
3 mM SAM	0.5 μl
[³ H]-SAM (67.3 Ci/mmol, 0.5 μCi/μl)	1 μl
RNase inhibitor 40 U/μl	1 μl
RNase free water	up to 30 μl
Total	30 μl
4. Incubate the reaction at 30-37 °C depending on different MTases for 1.5 h.
5. Transfer the tubes onto ice and add equal volume (30 μl) of 0.2% SDS, 20 mM EDTA.
6. Keep the tubes on ice, add 1 ml of 10 mM NH₄HCO₃ (pH 8.5).
7. Prepare 1 ml DEAE Sephadex column and equilibrated with 10 ml 10 mM NH₄HCO₃ (pH 8.5).
8. Load the samples onto the equilibrated column.
9. Wash the column with 10 ml of 10 mM NH₄HCO₃ (pH 8.5), 100 mM NaCl.
10. Elute the samples with 1.5 ml of 10 mM NH₄HCO₃ (pH 8.5), 400 mM NaCl.
11. Add equal volume scintillation liquid, mix well by vortexing and measure the signal using Liquid scintillation counter. The counting signal of [³H], which is transformed from [³H]-SAM to RNA substrates, represents the activity of tested MTases.

Recipes

1. 10x MTase Buffer
 - 0.5 M Tris-HCl (pH 7.5 or 8.0)
 - 50 mM KCl
 - 20 mM MgCl₂
 - 20 mM DTT
2. Cap-0 cap structure (m7GpppN-RNA)

Heat denatured RNA	13.5 μ l
10x Bodicon Capping Buffer	2 μ l
10 mM GTP	2 μ l
3 mM SAM	1 μ l
RNase inhibitor 40 U/ μ l	0.5 μ l
Bodicon Capping Enzyme (10 U/ μ l)	2 μ l
Total	20 μ l
3. Non-methylated Cap-0 cap structure (GpppN-RNA)	
Heat denatured RNA	13.5 μ l
10x Bodicon Capping Buffer	2 μ l
10 mM GTP	2 μ l
Inorganic pyrophosphatase 0.1 U/ μ l	1 μ l
RNase inhibitor 40 U/ μ l	0.5 μ l
Bodicon Capping Enzyme (10 U/ μ l)	2 μ l
Total	20 μ l
4. MTase assay reaction mix	
Purified MTases	1 μ g
10x MTase Buffer	3 μ l
GpppN-RNA or m7GpppN-RNA	2 μ g
3 mM SAM	0.5 μ l
[³ H]-SAM (67.3 Ci/mmol, 0.5 μ Ci/ μ l)	1 μ l
RNase inhibitor 40 U/ μ l	1 μ l
RNase free water	up to 30 μ l
Total	30 μ l

Acknowledgments

We thank Dr. Tero Ahola from University of Helsinki for the kindly help and advices for setting up this MTase assay. We thankfully acknowledge the University of Helsinki for providing research facilities and support during our visits. This work was supported by the China NSFC grants (81130083, 31170152 and 81271817).

References

1. Chen, Y., Tao, J., Sun, Y., Wu, A., Su, C., Gao, G., Cai, H., Qiu, S., Wu, Y., Ahola, T. and Guo, D. (2013). [Structure-function analysis of severe acute respiratory syndrome coronavirus RNA cap guanine-N7-methyltransferase](#). *J Virol* 87(11): 6296-6305.
2. Chen, Y., Su, C., Ke, M., Jin, X., Xu, L., Zhang, Z., Wu, A., Sun, Y., Yang, Z., Tien, P., Ahola, T., Liang, Y., Liu, X. and Guo, D. (2011). [Biochemical and structural insights](#)

[into the mechanisms of SARS coronavirus RNA ribose 2'-O-methylation by nsp16/nsp10 protein complex.](#) *PLoS Pathog* 7(10): e1002294.

3. Chen, Y., Cai, H., Pan, J., Xiang, N., Tien, P., Ahola, T. and Guo, D. (2009). [Functional screen reveals SARS coronavirus nonstructural protein nsp14 as a novel cap N7 methyltransferase.](#) *Proc Natl Acad Sci U S A* 106(9): 3484-3489.