

## Proteasome Assay in Cell Lysates

Pamela Maher\*

Cellular Neurobiology, Salk Institute for Biological Studies, La Jolla, USA

\*For correspondence: [pmaher@salk.edu](mailto:pmaher@salk.edu)

**[Abstract]** The ubiquitin-proteasome system (UPS) mediates the majority of the proteolysis seen in the cytoplasm and nucleus of mammalian cells. As such it plays an important role in the regulation of a variety of physiological and pathophysiological processes including tumorigenesis, inflammation and cell death (Ciechanover, 2005; Kisselev and Goldberg, 2001). A number of recent studies have shown that proteasome activity is decreased in a variety of neurological disorders including Parkinson's disease, Alzheimer's disease, amyotrophic lateral sclerosis and stroke as well as during normal aging (Chung *et al.*, 2001; Ciechanover and Brundin, 2003; Betarbet *et al.*, 2005). This decrease in proteasome activity is thought to play a critical role in the accumulation of abnormal and oxidized proteins. Protein clearance by the UPS involves two sequential reactions. The first is the tagging of protein lysine residues with ubiquitin (Ub) and the second is the subsequent degradation of the tagged proteins by the proteasome. We herein describe an assay for the second of these two reactions (Valera *et al.*, 2013). This assay uses fluorogenic substrates for each of the three activities of the proteasome: chymotrypsin-like activity, trypsin-like activity and caspase-like activity. Cleavage of the fluorophore from the substrate by the proteasome results in fluorescence that can be detected with a fluorescent plate reader.

### **Materials and Reagents**

1. Cells
2. HEPES (Sigma-Aldrich, catalog number: H3375)
3. MgCl<sub>2</sub> (Sigma-Aldrich, catalog number: M2670)
4. EDTA (Sigma-Aldrich, catalog number: E5134)
5. EGTA (Sigma-Aldrich, catalog number: E4378)
6. Sucrose (MP Biomedicals, catalog number: 821713)
7. DTT (Life Technologies, catalog number: 15508013)
8. Proteasome substrates [dissolved in DMSO (Sigma-Aldrich, catalog number: D8418) to a final concentration of 10 mM and stored frozen at -20 °C]
  - a. Suc-LLVY-AMC (chymotrypsin-like activity substrate) (Enzo Life Sciences, catalog number: P802)
  - b. Z-ARR-AMC (trypsin-like activity substrate) (EMD Millipore, catalog number: 539149)

- c. Z-LLE-AMC (caspase-like activity substrate) (EMD Millipore, catalog number: 539141)
9. PBS without  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  (Sigma-Aldrich, catalog number: P802)
10. Coomassie Protein Assay Reagent (Thermo Fisher Scientific, catalog number: 1856209)
11. BSA protein standard (Thermo Scientific, catalog number: 23209)
12. ATP (Sigma-Aldrich, catalog number: A3377)
13. Proteasome Lysis/Assay Buffer (see Recipes)

### **Equipment**

1. 60 mm tissue culture dishes
2. Rubber policeman or other type of cell scraper
3. 1.7 ml microcentrifuge tube
4. Black walled 96 well plates (Corning, Costar<sup>®</sup>, catalog number: 3603)
5. Clear 96 well plates (Greiner Bio-One GmbH, catalog number: 655101)
6. Sonicator (GlobalSpec, model: W-380)
7. Microcentrifuge
8. Fluorescent plate reader
9. Visible plate reader

### **Procedure**

#### **A. Sample Preparation**

1. Rinse cells in 60 mm dishes 2 times with ice cold PBS.
2. Scrape cells using a rubber policeman or other type of cell scraper into 400  $\mu\text{l}$  lysis buffer, transfer to a 1.7 ml microcentrifuge tube and place on ice.
3. Sonicate for 10 sec using microtip set on ~2.
4. Centrifuge at 16,000  $\times g$  for 10 min at 4 °C.
5. Transfer supernatant to a fresh 1.7 ml microcentrifuge tube (can store at -70 °C until use in assay).

#### **B. Assay**

1. Prepare assay buffer (make up more than you need; e.g. if you need 3 ml, make 3.5 ml).
2. Add proteasome substrates to assay buffer (2.5  $\mu\text{l}$ /sample; 100  $\mu\text{M}$  final concentration) (make up slightly more than you need; e.g. if you have 12 samples, make up enough for 13).
3. Put 2  $\times$  200  $\mu\text{l}$  aliquots of assay buffer with substrate into 2 wells of a black walled 96 well plate for each sample.

4. Add 50  $\mu$ l cell lysate to each well (include 2 wells with lysis buffer with no cells as assay blanks).
5. Incubate 60 min at 37 °C in order to allow the cleavage of the fluorophore from the proteasome substrate.
6. Read  $A_{360\text{ex}}/A_{460\text{em}}$  on fluorescent plate reader.
7. Normalize to protein determined with the Coomassie Protein Assay following the manufacturer's instructions for a microplate reader and using 5  $\mu$ l of lysate.

### **Recipes**

1. Proteasome Lysis/Assay Buffer

50 mM HEPES (pH 7.8)

10 mM NaCl

1.5 mM  $\text{MgCl}_2$

1 mM EDTA

1 mM EGTA

250 mM sucrose

Sterile filter and stored at 4 °C

For lysis buffer: add DTT to 5 mM final concentration (use 1 M stock)

For assay buffer: add DTT to 5 mM final concentration and ATP to 2 mM final concentration

### **Acknowledgments**

This protocol was adapted from Valera *et al.* (2013). The research was supported by grants from NIH (RO1AG035055), the Fritz B. Burns Foundation and the Alzheimer's Association.

### **References**

1. Betarbet, R., Sherer, T. B. and Greenamyre, J. T. (2005). [Ubiquitin-proteasome system and Parkinson's diseases](#). *Exp Neurol* 191 Suppl 1: S17-27.
2. Chung, K. K., Dawson, V. L. and Dawson, T. M. (2001). [The role of the ubiquitin-proteasomal pathway in Parkinson's disease and other neurodegenerative disorders](#). *Trends Neurosci* 24(11 Suppl): S7-14.
3. Ciechanover, A. (2005). [Proteolysis: from the lysosome to ubiquitin and the proteasome](#). *Nat Rev Mol Cell Biol* 6(1): 79-87.

4. Ciechanover, A. and Brundin, P. (2003). [The ubiquitin proteasome system in neurodegenerative diseases: sometimes the chicken, sometimes the egg.](#) *Neuron* 40(2): 427-446.
5. Kisselev, A. F. and Goldberg, A. L. (2001). [Proteasome inhibitors: from research tools to drug candidates.](#) *Chem Biol* 8(8): 739-758.
6. Valera, E., Dargusch, R., Maher, P. A. and Schubert, D. (2013). [Modulation of 5-lipoxygenase in proteotoxicity and Alzheimer's disease.](#) *J Neurosci* 33(25): 10512-10525.