

Fluorescence Microscopy for Cilia in Cultured Cells and Zebrafish Embryos

Jingli Cao, Xueliang Zhu and Xiumin Yan*

State Key Laboratory of Cell Biology, Institute of Biochemistry and Cell Biology, Shanghai, China

*For correspondence: yanx@sibcb.ac.cn

[Abstract] Cilia are microtubule-based hair-like projections found in organisms, ranging from protozoa to mammals. This protocol provides methods for immunofluorescence staining of cilia in cultured cells and zebrafish embryos.

Materials and Reagents

1. hTERT-RPE1 cell line (ATCC , catalog number: CRL-4000)
2. Zebrafish (AB strain) (China Zebrafish Resource Center)
3. DMEM/F12 medium (Life Technologies, Invitrogen™, catalog number: 11330-032)
4. Fetal bovine serum (Life Technologies, Invitrogen™, catalog number: 16000-044)
5. Hygromycin B (Life Technologies, Invitrogen™, catalog number: 10687010)
6. Paraformaldehyde (Sigma-Aldrich, catalog number: P6148)
7. Pure ethanol (Sinopharm Chemical Regent, catalog number: 10009218)
8. Low-melting-point agarose (Sigma-Aldrich, catalog number: A9414)
9. Bovine Serum Albumin (BSA) (Sigma-Aldrich, catalog number: A3912)
10. Triton X-100 (AMRESCO, catalog number: 0694)
11. Tween-20 (AMRESCO, catalog number: 0777)
12. Mouse monoclonal anti-acetylated tubulin antibody (Sigma-Aldrich, catalog number: T-6793)
13. Goat anti-mouse IgG (H+L) antibody conjugated with Alexa Fluor-546 (Life Technologies, catalog number: A11030)
14. Donkey anti-mouse IgG (H+L) antibody conjugated with Alexa Fluor-488 (Life Technologies, catalog number: A21202)
15. 4,6-diamidino-2-phenylindole (DAPI) (Sigma-Aldrich, catalog number: D8417)
16. Dako mounting medium (Dako, catalog number: s3023)
17. PBS (see Recipes)
18. PBT (see Recipes)
19. 4% paraformaldehyde (PFA) fixation solution (see Recipes)
20. Embryo fixation buffer (see Recipes)
21. Embryo blocking buffer (see Recipes)

22. 4', 6-diamidino-2-phenylindole (DAPI) stock solution (see Recipes)
23. Complete growth medium (see Recipes)
24. Serum free medium (see Recipes)
25. 75% ethanol (see Recipes)

Equipment

1. Millex-GP Filter unit with 0.22 µm pore size (Millipore, catalog number: SLGP033RS)
2. Forceps (The duMONT Company, catalog number: 0203-5/15-PO)
3. 10 cm Petri dishes (BD Biosciences, Falcon[®], catalog number: 353003)
4. 12-well plates (Corning, catalog number: 3513)
5. 2 ml Eppendorf (EP) tubes (Axygen, catalog number: MCT-200-C)
6. Glass slides (Fan Yi, catalog number: 7105P)
7. 18mm diameter circle cover slips (Thermo Fisher Scientific, catalog number: 12-545-84)
8. Parafilm (Parafilm M[®], catalog number: PM-996)
9. 100 ml beaker (Thermo Fisher Scientific, catalog number: 1201-0100)
10. Filter paper (GE, catalog number: 10311611)
11. Plastic transfer pipette (Thermo Fisher Scientific, catalog number: 11387873)
12. Shakers (Qilinbeier, catalog number: TS-8S)
13. Leica TCS SP5 confocal microscope equipped with a 63x oil objective and a 63x NA1.2 water immersion objective

Procedure

A. Immunofluorescence labeling of cilia in cultured cells

1. hTERT-RPE1 cells are cultured as described in the ATCC website (<http://www.atcc.org/Products/All/CRL-4000.aspx#culturemethod>). Briefly, cells are grown in 10 cm Petri dishes with 10-12 ml complete growth medium (see Recipes) and media are changed every other day. Cells are subcultivated at 1:5 ratio twice a week when cell concentration reaches between 2×10^4 and 4×10^4 cells/cm². An inoculum of 4×10^3 to 6×10^3 viable cells/cm² is recommended.
2. Sterilize 18 mm diameter circle cover slips by immersing in 75% ethanol in a 100 ml beaker over night. Pick cover slips by using forceps and put slips into 12-well-plate, one slip per well. Rinse slips three times with 1 ml PBS per well each time. Add 1 ml complete growth medium to each well, and then add appropriate aliquots of the dissociated cell suspension.

3. Culture for 1-2 days until cells reach 60-90% confluence. Wash three times with 1 ml PBS per well each time to get ride of serum, then add 1 ml serum free medium and culture for 48 h to induce cilia formation. The culture time and confluence should be determined by researchers themselves to get the best result for different experiments. Generally, a confluence higher than 60% is recommended for efficient cilia induction.
4. Before fixation, place cells on ice for 30 min to destabilize microtubules and rinse once with 1 ml ice-cold PBS for each well.
5. Fix cells with 1 ml of freshly prepared 4% PFA in PBS per well for 10 min at room temperature (RT).
6. Briefly rinse cells three times with 1 ml PBS each time per well, and permeabilize cells with 1 ml 0.5% TritonX-100 in PBS per well for 15 min at RT.
Note: PFA waste should be collected in an appropriate container for proper disposal.
7. Briefly rinse cells three times with 1 ml PBS each time per well.
8. Incubate cells with 1 ml 1% BSA in PBS each well for 1 h at RT.
9. Make a humid chamber with a new 10 cm Petri dish by putting a piece of wet filter paper and a piece of parafilm on the top of the paper. Make sure the parafilm is flat. Transfer the cover slips with blocked cells by using forceps onto the parafilm. Add 100 μ l diluted mouse anti-acetylated tubulin antibody (1:1,000 in PBS with 1% BSA) onto each slip and incubate for 2 h at RT or overnight at 4 °C.
10. Wash cells with 200 μ l 1% BSA in PBS per slip three times for 5 min each.
11. Add 100 μ l diluted secondary antibody, Goat anti-mouse IgG (H+L) antibody conjugated with Alexa Fluor-546 (1:1,000 in PBS with 1% BSA) for 1 h at RT in the dark.
12. Wash cells with 200 μ l 1% BSA in PBS per slip three times for 5 min each.
13. To stain nuclei, dilute DAPI stock solution with PBS to 0.5 μ g/ml, add 100 μ l diluted DAPI onto each slip and incubate for 1 min at RT.
14. Wash cells with 200 μ l PBS per slip three times for 5 min each.
15. Mount slides with 10-20 μ l Dako mounting medium as shown in Figure 1 and dry them for at least 2 h at RT in the dark. Store slides at 4 °C. Samples are good for at least 1 month, but DAPI signals will diffuse and become dim after 1 week. It's better to take images within 1 week.
16. Take images on a Leica TCS SP5 confocal microscope with a 63x oil objective. Optical sections are captured at 0.5 μ m intervals and z-stack images are obtained by maximum intensity projections. A typical image can be found in Figure 1j of Cao *et al.* (2012).

B. Whole mount immunofluorescence of zebrafish embryos

1. Zebrafish (AB strain) were raised and maintained under standard conditions and staged as previously described in hours post fertilization (hpf) (please refer to *The Zebrafish Book*: http://zfin.org/zf_info/zfbook/zfbk.html).
2. Dechorionate zebrafish embryos at the 8-somite stage or at the 24 hpf stage through pronase treatment or manually with forceps. For detailed method about dechoriation, please refer to Chapter 4 of *The Zebrafish Book* (http://zfin.org/zf_info/zfbook/chapt4/4.1.html).
3. Transfer embryos with plastic transfer pipette into 2 ml EP tubes, up to 100 embryos each tube. Remove extra water and fix embryos with 1 ml embryo fixation buffer for 2 h at RT or overnight at 4 °C with gentle rocking at 15-20 times per minute on a shaker. All washings and incubations should be done with gentle rocking. All solutions are added and removed by using pipette with 1 ml tips. PFA and methanol waste should be collected in an appropriate container for proper disposal.
4. After a brief rinse with 1 ml PBT, dehydrate embryos through a graded methanol/PBT series (25%, 50%, 75%, 100%). Rock the embryos gently at RT for 5 min at each step.
5. Incubate embryos overnight in 1ml 100% methanol at -20 °C. Embryos can be stored at -20 °C for up to 1 month before further staining.
6. Rehydrate embryos in a methanol/PBT graded series (75%, 50%, 25%, 5 min each at RT). Rinse three times with PBT for 5 min each at RT, and use 1 ml PBT each time per tube.
7. Incubate embryos for 1-2 h in 1 ml embryo blocking buffer at RT.
8. Incubate samples overnight at 4 °C with mouse anti-acetylated tubulin antibody diluted in 300-500 µl blocking buffer (1:1,000).
Note: At least 300 µl is recommended for sufficient incubation.
9. Wash three times for 10 min each, with 1ml 0.5% Triton X-100 in PBS each time.
10. Incubate embryos with 300-500 µl secondary antibody, Donkey anti-mouse IgG (H+L) antibody conjugated with Alexa Fluor-488 diluted in blocking buffer (1:1,000) overnight at 4 °C or for 2 h at RT.
11. Rinse three times for 10 min each with 1ml 0.5% Triton X-100 in PBS each time.
12. To stain nuclei, dilute DAPI stock solution with PBT to 0.5 µg/ml, and incubate embryos with 1 ml diluted DAPI each tube for 5-10 min at RT. Then wash embryos three times for 5 min each with 1 ml PBT. Embryos are now ready for embedding. Otherwise, please keep stained embryos in PBT at 4 °C for up to 1 week.
13. Embed embryos in 1% low melting agarose. Briefly, prepare a 1% low-melting-point agarose solution in H₂O by heating in a microwave oven. Keep the solution liquid in a tube in a 37 °C water bath. This solution remains liquid at 37 °C, but quickly hardens when cooled lower than 30 °C. Use 35 mm culture dishes for embryos mounting and

imaging. First, add 1 ml agarose solution in each dish and let it gel at RT. Make sure the agarose is flat and covers the whole bottom of the dish. Transfer four embryos by using plastic transfer pipette onto the agarose gel of each dish. Quickly pipette a small drop of agarose onto the embryo, and position the embryo at the appropriate angle in the agarose using forceps and allow the agarose to harden. Make sure the embryos are distant from each other, so that you can mount one by one. 24 hpf embryos should be mounted laterally to show the pronephric duct, and 8-somite-stage embryos should be mounted to show the Kupffer's Vesicle on the up side. Please refer to Figure 7a and b of the Reference 1 for correct orientations. Mounted embryos should be examined at the same day.

14. Add 2-3 ml H₂O into the 35 mm dishes with mounted embryos, and take images on a Leica TCS SP5 MP confocal microscope with a 63x NA1.2 water immersion objective. Optical sections are captured at 0.5- μ m intervals and z-stack images are obtained by maximum intensity projections. A typical image can be found in Figure 7g and j of Cao *et al.* (2012).




Figure 1. Mounting cells for microscopy

Recipes

1. PBS
 - 137 mM NaCl
 - 2.7 mM KCl
 - 8 mM Na₂HPO₄

- 2 mM KH₂PO₄
pH 7.4
Autoclaved for long time storage
2. PBT
PBS with 0.1% Tween-20
 3. 4% PFA
Dissolve 4 g PFA in 100 ml PBS, heat to 60 °C for 1 h with stirring
The cooled solution can be filtered through Millex-GP Filter unit with 0.22µm pore size, and stored at -20 °C. It is good for a month at -20 °C, or use immediately after preparation, which is recommended.
 4. Embryo fixation buffer
4% PFA in PBS supplemented with 0.5% Triton X-100
 5. Embryo blocking buffer
2% BSA
0.5% normal goat serum
1% DMSO
0.5% Triton X-100 in PBS
 6. 4', 6-diamidino-2-phenylindole (DAPI) stock solution
Dissolve DAPI powder in dimethylformamide (DMF) to make a 5 mg/ml stock solution
 7. Complete growth medium
DMEM/F12 medium supplemented with 10% fetal bovine serum and 0.01 mg/ml hygromycin B
 8. Serum free medium
DMEM/F12 medium supplemented with 0.01 mg/ml hygromycin B
 9. 75% ethanol
75 ml pure ethanol mixed with 25 ml H₂O

Acknowledgments

This protocol was adapted from previous work (Cao *et al.*, 2012; Chapter 4 of The Zebrafish Book (http://zfin.org/zf_info/zfbook/chapt4/4.1.html)). The work was supported by the National Basic Research Program of China (2012CB945003 and 2010CB912102), National Science Foundation of China (30971430, 30830060, and 31010103910), and Chinese Academy of Sciences (XDA01010107).

References

1. Cao, J., Shen, Y., Zhu, L., Xu, Y., Zhou, Y., Wu, Z., Li, Y., Yan, X. and Zhu, X. (2012). [miR-129-3p controls cilia assembly by regulating CP110 and actin dynamics](#). *Nat Cell Biol* 14(7): 697-706.