

Preparation of Golgi Membranes from Rat Liver

Carmen Valente, Gabriele Turacchio, Stefania Spanò, Alberto Luini and Daniela Corda*

Institute of Protein Biochemistry (CNR), Naples, Italy

*For correspondence: d.corda@ibp.cnr.it

[Abstract] This protocol details the isolation of enriched Golgi membranes from rat liver, using discontinuous density gradient centrifugation. This high-yield extraction method is useful for several applications, including immunoprecipitation of solubilised Golgi membrane proteins (preparation included) and electron microscopy. Protocol adapted from Leelavathi *et al.* (1970).

Materials and Reagents

1. Fresh rat liver tissue (30 mg)
2. HEPES (Sigma-Aldrich, catalog number: H4034)
3. KCl (Baker, catalog number: 0208)
4. Triton X-100 (ACROS, catalog number: 215680010)
5. EDTA (Baker, catalog number: 1073)
6. Sodium orthovanadate (Sigma-Aldrich, catalog number: S6508)
7. β -Glycerophosphate (Sigma-Aldrich, catalog number: G9422)
8. NaF (Sigma-Aldrich, catalog number: S7920)
9. Protease inhibitor cocktail (complete Mini EDTA-free) (Roche Diagnostics; or equivalent) (Roche Diagnostics, catalog number: 05056489001)
10. Protein assay kit (Bio-Rad Laboratories, catalog number: 500-0006)
11. $MgCl_2$ (Sigma-Aldrich, catalog number: 16104)
12. Sucrose (Sigma-Aldrich, catalog number: 16104; extra pure grade) (see Recipes)
13. Phosphatase inhibitor cocktail (Roche Diagnostics, catalog number: 04906837001) (see Recipes)

Equipment

1. Ultraturrax homogeniser (T 25 digital ULTRA-TURRAX® IKA® Werke Staufen/Germany)
2. Centrifuge (Ultracentrifuge, Beckman Coulter, model: Optima XE-100; ultra clear centrifuge tubes, catalog number: 344058)
3. SW28 (or equivalent) swing-out rotor [swinging bucket rotor, catalog number: 2352; Beckman SW 28 (Beckman Coulter, catalog numbers: 342207, 342204), with titanium

- buckets 6 x 38.5 ml, 25 x 89 mm, 28,000 rpm, 141,000 x g]
4. Bausch and Lomb refractometer (Lab Logistics Group GmbH)

Procedure

1. About 30 mg rat liver tissue is excised from six male Sprague-Dawley rats (weighing 200-300 g each) fed *ad libitum* the day before the Golgi-membrane preparation. The tissue is placed into 100 ml ice-cold 0.5 M sucrose solution and squeezed to expel the blood. Then, the tissue is minced to a fine purée with scissors in 140 ml ice-cold 0.5 M sucrose solution, and then homogenised for 30 sec at 12 Watt using an Ultraturrax homogeniser (or equivalent) at 4 °C.
2. The homogenate is centrifuged for 10 min at 600 x g at 4 °C.
3. The supernatant (~120 ml in 0.5 M ice-cold sucrose solution) is then split into 20 ml aliquots, each of which is slowly layered onto an 18 ml 1.3 M sucrose cushion by being carefully dripped down the side of the 38.5 ml centrifuge tubes (so as not to disturb the cushion) for the SW28 (or equivalent) swing-out rotor. Centrifuge the cushion plus homogenate supernatant for 60 min at 105,000 x g at 4 °C in an SW28 swing-out rotor. This results in the formation of: (a) an upper fat layer; (b) a clear reddish supernatant; (c) a thick membrane felt layer above the 1.3 M sucrose interface; (d) a turbid 1.3 M sucrose supernatant; and (e) a sediment at the bottom of the tube.
4. The interface between the 1.3 M and 0.5 M sucrose layers (~2.5 ml per tube) is collected and used for the preparation of the Golgi-rich fraction. This is collected by carefully puncturing the tube at the middle of the interface with an 18 gauge needle (3.8 cm long) and using a syringe to extract the interface. This is gently resuspended and the molarity of the sucrose adjusted to 1.1 M by adding 2.3 M sucrose (usually to a final volume of ~16 ml), with the aid of a Bausch and Lomb (or equivalent) refractometer. The volume is then brought to a total of 18 ml with 1.1 M sucrose.
5. The suspension is split into two 9-ml aliquots that are layered onto sucrose gradients that are previously prepared using 7.5 ml each of 1.4 M, 1.3 M and then 1.25 M sucrose, layered on top of each other from the bottom of the cellulose acetate tubes (using the same tubes as in step 3) for the SW28 (or equivalent) swing-out rotor. Finally, the 1.1 M sucrose sample layer is covered with 6.5 ml 0.5 M sucrose solution. These stages are all carried out with care taken not to disturb or mix the layers of the gradient in any way. Prior to the layering of the sample onto the 1.25 M sucrose, the prepared gradients are stored for at least 20 min at 4 °C.
6. These are centrifuged for 90 min at 105,000 x g at 4 °C, and then the Golgi-enriched membrane fraction is collected at the interface between the 0.5 M and 1.1 M sucrose (as

- for step 4).
7. The collected Golgi-enriched membrane fractions are pooled and assayed for protein concentration using a commercially available protein assay kit, according to the manufacturer instructions. This provides an estimate of the yield of the Golgi fraction and its contamination with rough endoplasmic reticulum membranes and/or free ribosomes. The yield for the Golgi-enriched fraction is about 1.5 mg to 2.0 mg of protein per 1 g liver. The pooled Golgi-enriched membrane fractions are then aliquoted as required, frozen in liquid nitrogen, and stored at -80 °C. Routinely, for batch control variations, Western blotting analysis of the Golgi-enriched fraction protein is performed using antibodies against Golgi-localised and/or endoplasmic reticulum-localised proteins (e.g., anti-mannosidase and anti-protein disulphide isomerase antibodies, respectively).
 8. To solubilise these Golgi-enriched membranes for use in pull-down assays, 4 mg Golgi-enriched membrane proteins (800 µl) are diluted two-fold (1:1) in 10 mM HEPES (pH 8.0), 50 mM KCl, with protease and phosphatase inhibitor cocktails added, centrifuged (10 min, 20,000 x g, 4 °C). This wash step is then repeated again, resuspending the Golgi-enriched membrane pellets in 800 µl of the same buffer.
 9. The final Golgi-enriched membrane pellet is resuspended in 400 µl 20 mM HEPES (pH 8.0), 100 mM KCl, 1% Triton X-100, 2 mM EDTA, with protease and phosphatase inhibitor cocktails added, and solubilised for 30 min at 4 °C.
 10. The solubilised suspension is then diluted 1:1 with water (to a final 800 µl volume), and centrifuged for 10 min at 2,000 x g at 4 °C. The supernatant from this final centrifugation is used as the “solubilised Golgi-enriched membranes”.

Recipes

1. Sucrose
 Sucrose solutions (0.5 M, 1.1 M, 1.25 M, 1.3 M, 1.4 M, 2.3 M) in 0.1 M potassium phosphate buffer (pH 6.65) (combine 38.1 ml of 1 M K₂HPO₄ stock solution with 61.9 ml of 1 M KH₂PO₄ stock solution and then dilute to 1 L with distilled H₂O) with 5 mM MgCl₂
2. Phosphatase inhibitor cocktail
 1 µM sodium orthovanadate
 20 mM β-glycerophosphate
 10 mM NaF
 This cocktail is made up in water at 4 °C, immediately before use.

Acknowledgments

The authors would like to thank: The Italian Association for Cancer Research (to D. C. IG4664, IG10341; to A. L. IG4700), Telethon Italia (to D. C. GGPO9274; to A. L. GGPO8231), the European Community Seventh Framework Programme FP7/2007-2013 HEALTH-F2-2007-201804 (Eucilia to A. L.), grant FIT DM 24/09/2009, Legge 46/82, Ministry of Economy and Finance (to D. C.), PON project 01/00117 2007-2013 “Antigens and adjuvants for vaccines and immunotherapy” and the MIUR Project “FaReBio di Qualità”, for financial support. C. V. and S. S. were recipients of Italian Foundation for Cancer Research (FIRC, Milan, Italy) Fellowships. This protocol was adapted from Leelavathi *et al.* (1970), as detailed in Valente *et al.* (2012). The authors also thank C.P. Berrie for editorial assistance.

References

1. Leelavathi, D., Estes, L., Feingold, D. S. and Lombardi, B. (1970). [Isolation of a Golgi-rich fraction from rat liver](#). *Biochim Biophys Acta* 211(2): 124-138.
2. Valente, C., Turacchio, G., Mariggio, S., Pagliuso, A., Gaibisso, R., Di Tullio, G., Santoro, M., Formiggini, F., Spano, S., Piccini, D., Polishchuk, R. S., Colanzi, A., Luini, A. and Corda, D. (2012). [A 14-3-3gamma dimer-based scaffold bridges CtBP1-S/BARS to PI\(4\)KIIIbeta to regulate post-Golgi carrier formation](#). *Nat Cell Biol* 14(4): 343-354.