

Protocol for Whole Cell Lysis of Yeast

Zongtian Tong*

Department of Cell Biology, Center for Metabolism and Obesity Research, Johns Hopkins School of Medicine, Baltimore, USA

*For correspondence: tongzong@gmail.com

[Abstract] This protocol describes how to perform lysis on whole yeast cell samples using NaOH. The lysed cells can then be used for downstream applications such as the extraction of total proteins.

Materials and Reagents

1. NaOH (Sigma-Aldrich)
2. Beta-mercaptoethanol (beta-ME) (Sigma-Aldrich)
3. 100% (w/v) trichloroacetic acid (TCA) (Sigma-Aldrich)
4. Pepstatin A (Sigma-Aldrich, catalog number: P5318)
5. Leupeptin (Sigma-Aldrich, catalog number: L2884)
6. PMSF (Sigma-Aldrich, catalog number: P7626)
7. Acetone (Sigma-Aldrich)
8. Protease inhibitors
9. DMSO
10. Glycerol
11. BPB
12. Ethanol
13. SDS lysis buffer
14. 7.5 ml 1 M Tris-HCl (pH 6.8) (see Recipes)
15. DTT stock (see Recipes)
16. 5x SDS-PAGE gel loading dye (see Recipes)
17. Protease inhibitors (see Recipes)

Procedure

1. Pellet 2 OD of cells. Wash 1x with 0.5 ml of water (cell pellets may be frozen in liquid nitrogen at this point).
2. Resuspend pellets in 100 μ l of water (make sure pellets are fully resuspended).

3. Add 17 μ l of 1.85 M *NaOH/7.4% (v/v) beta-ME and vortex. Incubate on ice for 10 min.
4. Add 8 μ l of 100% (w/v) TCA, vortex, and incubate on ice for 10 min.
5. Spin extract for 10 min at 20,000 \times g /14,000 rpm at 4 °C. Wash protein pellet with 500 μ l of cold acetone (do not pipette up and down; add the solution and vortex; it is OK if the pellet is not completely re-suspended).
6. Re-spin sample for 5 min at 4 °C. Dry almost completely. If the pellet gets too dry, it will not dissolve well later.
7. Resuspend pellet in 100 μ l of SDS lysis buffer + 1x protease inhibitors (pipette up and down, then vortex, then resuspend the pellet completely).
8. If doing a BCA assay, boil for 5 min and perform the assay.
9. Add 25 μ l of 5x SDS-PAGE gel loading dye to 100 μ l of sample.
10. Boil for 5 min and load 40 μ g total proteins per lane.

Recipes

1. To make 1 ml of NaOH/beta-ME, mix 741 μ l of water, 185 μ l of 10 N NaOH (stock 10 N NaOH solution) and 74 μ l of beta-ME stock. Always make fresh NaOH/beta-ME.
2. 5x SDS-PAGE gel loading dye
 - 150 mM Tris-HCl
 - 15% SDS
 - 25% glycerol
 - 0.02% BPB
 - 12.5% beta-ME
3. To make 7.5 ml 1 M Tris-HCl (pH 6.8)
 - 7.5 g SDS
 - 25 ml 50% glycerol
 - 10 ml H₂O
 - 20 mg BPB

Bring final volume to 43.75 ml. Aliquot into 1 ml and add 125 μ l of beta-ME fresh prior to use. SDS may come out of solution, but heating will fix this.
4. Protease Inhibitors
 - Leupeptin/pepstatin A (1 mg/ml pepstatin A dissolved in DMSO; 10 mg/ml leupeptin dissolved in ddH₂O)
 - Combine and aliquot 1 ml per tube and store at -20 °C.
 - PMSF: 0.5 M dissolved in ethanol. Store in 1 ml aliquots at -20 °C. Supplied as 1,000x stock.
5. DTT stock (1 M)

Dissolve in ddH₂O, store in 1 ml aliquots at -20 °C. *I.e.* 1,000x stock.

Acknowledgments

This protocol has been modified and adapted in the Espenshade Lab, Johns Hopkins School of Medicine. Funding to support different projects that have used this protocol has come from NIH – National Heart, Lung, and Blood Institute, National Institute of Allergy and Infectious Diseases, the Pancreatic Cancer Action Network, and the American Heart Association.

References

1. Horvath, A. and Riezman, H. (1994). [Rapid protein extraction from *Saccharomyces cerevisiae*](#). *Yeast* 10(10): 1305-1310.