

RNA Editing Detection by Direct Sequencing

Mehdi Jabnoue^{1*}, David Secco^{1,4}, Cécile Lecampion², Christophe Robaglia², Qingyao Shu³ and Yves Poirier¹

¹Department of Plant Molecular Biology, University of Lausanne, Lausanne, Switzerland; ²Laboratory of Plant Genetics and Biophysics, CEA Institute of Environmental Biology and Biotechnology, Aix Marseille University, Faculty of Sciences, Marseille, France; ³Institute of Nuclear Agricultural Sciences, Zhejiang University, Hangzhou, China; ⁴Australian Research Council Centre of Excellence in Plant Energy Biology, The University of Western Australia, Perth, Australia

*For correspondence: mehdi.jabnoue@gmail.com

[Abstract] RNA editing is a widespread post-transcriptional phenomenon through which primary RNA sequences are altered by nucleotide insertion/deletion or base conversion. It occurs in a variety of organisms and cooperates with alternative splicing in increasing both proteomic and transcriptomic complexity. We describe here a method allowing RNA editing events detection by performing direct sequencing of both genomic DNA and cDNA from the same source.

Materials and Reagents

1. DNeasy Plant mini kit (QIAGEN, catalog number: 69104)
2. RNeasy mini kit (QIAGEN, catalog number: 74104)
3. pJET PCR cloning system (Thermo Fisher Scientific, Fermentas, catalog number: K1231)
4. Recombinant DNase I (Life Technologies, Ambion[®], catalog number: AM2235)
5. M-MLV reverse transcriptase (Promega corporation, catalog number: 28025-013)
6. Oligo(dT) 15 Primer (500 mg) (Promega corporation, catalog number: C1101)
7. KOD high fidelity polymerase (TOYOBO, catalog number: KOD-201 200U)
8. Zymoclean Gel DNA Recovery Kit (ZYMO RESEARC, catalog number: D4001)

Equipment

1. NanoDrop (Thermo Fisher Scientific)

Procedure

1. DNA and total RNA were extracted with respectively DNeasy Plant mini kit and RNeasy mini kit, according to the manufacturer's instructions.
2. Each RNA sample was DNase treated (Recombinant DNase I) and quantified with a Nanodrop.
3. cDNAs were generated by M-MLV reverse transcriptase using oligo (dT) according to the manufacturer's instructions.
4. The cDNA produced was diluted with four parts of sterile deionized water.
5. PCR amplification of the target (containing the putative editing region) was performed using KOD high fidelity polymerase and target specific oligonucleotide primers (it's better to design primers that spans introns to prevent the amplification of residual contaminant genomic DNA).
6. PCR product was purified using a Zymoclean Gel DNA Recovery Kit and cloned into a pJET vector using pJET cloning kit.
7. Direct sequencing was performed on plasmid containing the target cloned in pJET vector. For each sample, 2 independent RT-PCR reactions were performed.

Acknowledgments

This work was funded by the Swiss National Foundation grant (31003A-12293 and 31003A-138339) and the Sino-Swiss Science and Technology Cooperation Program (IZLCZ3 123946 to YP and 2009DFA32040 to QS).

References

1. Eisenberg, E., Li, J. B. and Levanon, E. Y. (2010). [Sequence based identification of RNA editing sites](#). *RNA Biol* 7(2): 248-252.
2. Nakae, A., Tanaka, T., Miyake, K., Hase, M. and Mashimo, T. (2008). [Comparing methods of detection and quantitation of RNA editing of rat glycine receptor alpha3](#). *Int J Biol Sci* 4(6): 397-405.
3. Picardi, E., Gallo, A., Galeano, F., Tomaselli, S. and Pesole, G. (2012). [A novel computational strategy to identify A-to-I RNA editing sites by RNA-Seq data: de novo detection in human spinal cord tissue](#). *PLoS One* 7(9): e44184.
4. Ramaswami, G., Lin, W., Piskol, R., Tan, M. H., Davis, C. and Li, J. B. (2012). [Accurate identification of human Alu and non-Alu RNA editing sites](#). *Nat Methods* 9(6): 579-581.