

Mismatched Primer Extension Assays

Vasudevan Achuthan and Jeffrey J. DeStefano*

Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, USA

*For correspondence: jdestefa@umd.edu

[Abstract] Steady state kinetic assays have been a reliable way to estimate fidelity of several polymerases (Menendez-Arias, 2009; Rezende and Prasad, 2004; Svarovskaia *et al.*, 2003). The ability to analyze the extension of primers with specific mismatches at the 3' end is a major strength of the mismatched primer extension assays. Recently, we used the mismatched primer extension assays to show that the fidelity of HIV RT increases dramatically when concentration of Mg^{2+} is reduced to a physiologically relevant concentration (~0.25 mM) (Achuthan *et al.*, 2014). Here, we describe in detail how to perform the mismatched primer extension assay to measure the standard extension efficiency using human immunodeficiency virus reverse transcriptase (HIV RT) at 2 mM Mg^{2+} . The relative fidelity of the polymerase can then be estimated using the standard extension efficiency. The assay described here is based on the method published in Mendelman *et al.* (1990).

Materials and Reagents

1. Deoxynucleoside triphosphate (Roche Diagnostics, catalog number: 11969064001)
2. Gamma [γ - ^{32}P] ATP (PerkinElmer, catalog number: Blu502A001MC)
3. G-25 Macro spin columns (best suited for volumes of 75-150 μ l) (Harvard Apparatus, catalog number: 74-3901)
4. 40% Acrylamide-Bisacrylamide (19:1) solution (VWR International, catalog number: JT4969-0)
5. T4 polynucleotide kinase (PNK) (New England Biolabs, catalog number: M0201L)
6. 10X T4 polynucleotide kinase buffer (New England Biolabs, catalog number: B0201S)
7. Urea (VWR International, catalog number: 97061-926)
8. Ammonium Persulfate (VWR International, catalog number: 97064-594)
9. HIV Reverse Transcriptase, purified as described in Hou *et al.* (2004)
10. DNA oligonucleotides from Integrated DNA Technologies
 - a. Template:
5'-GGGCGAATTTAG[G/C]TTTTGTTCCCTTTAGTGAGGGTTAATTTTCGAGCTTG
G-3'. The underlined nucleotides in brackets indicate that templates with either a G or C at this position can be used depending on the type of mismatch examined.
 - b. Primer:

5'-TAACCCTCACTAAAGGGAACAAAAX-3'. "X" at the 3' end of the primer denotes A, T, or C depending on the mismatch examined. "X" in the case of a matched primer is G.

11. 1 M MgCl₂
12. Extension reaction buffer (see Recipes)
13. 2x loading dye (see Recipes)

Equipment

1. Eppendorf tubes
2. Micropipette
3. Table top centrifuge
4. Incubator
5. Gel apparatus

Software

1. Sigmaplot Version 10.0 (Sysstat Software)

Procedure

1. Primer labelling
 - a. All the primers should be first radiolabelled in 50 µl of 1x PNK buffer along with 50 pico moles of each primer, 10 µl of [γ -³²P] ATP and 5 units of PNK.
Note: The reaction mixture was incubated for 30 min at 37 °C and the PNK was heat inactivated for 15 min at 65 °C.
 - b. G-25 spin columns were incubated with 500 µl dH₂O for 15 min to equilibrate the column and the water was removed by spinning the columns at a table top centrifuge at 5,000 rpm for 4 min.
 - c. After heat inactivation, the excess [γ -³²P] ATP was removed from the reaction mixture by loading it onto an equilibrated column and spinning at 5,000 rpm for 4 min.
2. Matched primer extension reactions

To obtain information about the standard extension efficiency, extension of matched as well as mismatched primers should be performed. The standard extension efficiency can then be calculated as the ratio of efficiency of extending mismatched primers to efficiency of extending matched primers.

 - a. Eight matched primer extension reactions were set up. For each reaction, 14 nM of the radiolabeled primer was hybridized to 14 nM of the template (1:1 ratio of

primer:template) in 7 μ l of the extension reaction buffer. The mixture was heated at 65 °C for 5 min and then slowly cooled to room temperature.

- b. The hybrid was then incubated for 3 min at 37 °C in the reaction buffer along with 2 μ l of 10 mM MgCl₂ (final concentration of 2 mM MgCl₂) and 2 μ l of the nucleotide substrate (concentration varies for each reaction, see below) for each reactions. The nucleotide substrate is the next correct nucleotide to be added and it depends on the template used in the reactions. For this particular template, dCTP was the substrate (Figure 1). For matched primer extension reactions, the eight reactions had a final concentration of dCTP in the order of 0, 0.02, 0.04, 0.1, 0.2, 0.3, 0.6 and 1 μ M respectively.

Mismatched primer extension constructs:

XAAAACAAGGGAAATCACTCCCAAT-5'

5'-GGGCGAATTTAGCTTTTGTTCCCTTTAGTGAGGGTTAATTTTCGAGCTTGG-3'

X=A, T, G, or C

YAAAACAAGGGAAATCACTCCCAAT-5'

5'-GGGCGAATTTACGTTTTGTTCCCTTTAGTGAGGGTTAATTTTCGAGCTTGG-3'

Y=A, T, or C

Figure 1. Constructs used in mismatched primer extension assays. The sequence of the DNA templates is shown at the bottom and the sequence of the primer at the top. The underlined nucleotides show the only differences between the two templates.

- c. The extension was then initiated by addition of 2 μ l of 13 nM HIV RT (2 nM final concentration). The total reaction volume was 13 μ l.
- d. After 2 min, reactions were terminated by addition of 13 μ l of 2x loading dye.
Note: Reactions were run only for 2 min to ensure the primer is extended by only one nucleotide.
- e. The reaction products were then electrophoresed on 16% denaturing 7 M urea-polyacrylamide gels, dried, and imaged using a Fujifilm FLA5100 phosphorimager.
Note: The samples were run far enough to separate the extended band from the primer band (Figure 2).

3. Mismatched primer extension reactions

- a. For mismatched primer extension reactions, a different radiolabeled primer, depending on the mismatch analyzed (Figure 1), is used. Primer-template hybrids were made as described above.
- b. Eight individual reactions were set up. 7 μ l of primer-template hybrids was incubated at 37 °C in the reaction buffer for 3 min along with 2 μ l of 10 mM MgCl₂

(final concentration of 2 mM MgCl₂) and 2 μl of the nucleotide substrate. The total reaction volume was 13 μl.

Note: Mismatched primer-template sequences require more substrate for extension than matched primer-template sequences. So, the eight reactions had a final concentration of dCTP in the order of 0, 50, 100, 200, 400, 630, 1,200 and 1,870 μM respectively (Figure 2).

- c. Extension was initiated by addition of 2 μl of 13 nM HIV RT.
- d. After 5 min of extension, the reactions were terminated by addition of 13 μl of 2x loading dye and the extension products were processed on a 16% denaturing polyacrylamide gel as described above. The gel was run at 75 Watts for 90 min.




Figure 2. Representative data for the mismatched primer extension assay. Primer: template hybrid used here had a C.A mismatch at the 3' terminus. Concentration of dCTP used in each of the lanes (left to right): 0, 50, 100, 200, 400, 630, 1,200 and 1,870 μM.

4. Calculation of standard extension efficiency
 - a. Velocity measurements were performed according to Mendelman *et al.* (1990). Velocity (V_{rel}) of extending each primer by one nucleotide was calculated according to the formula:
$$V = (100 \times I_1) / \{ [I_0 + (0.5 \times I_1)] \times t \}$$
, where I_1 is the intensity of the extended band, I_0 is the intensity of the primer band, and t is the time of extension.
 - b. V_{rel} (percentage of primer extended per minute) was then plotted against [S], where [S] is the concentration of the substrate (dCTP) used in the reaction (Figure 3).

Note: These curves are typical of Michaelis-Menten kinetics curves.
 - c. The values of V_{max} and K_m were obtained by curve-fitting. The data were fitted to a “Single rectangular 2-parameter hyperbola equation” using the Sigmaplot software.

- d. The constant “a” obtained after curve fitting corresponds to the V_{max} and “b” corresponds to the K_m for this extension. V_{max} and K_m values were individually determined for both matched and mismatched primer extension reactions.
- e. The standard extension efficiency, f_{ext} , was calculated as the ratio of $\{V_{max}/K_m (mismatch)\}/\{V_{max}/K_m (match)\}$.

C.A mismatched primer extension kinetics




Figure 3. Representative graph for the mismatched primer extension assay. Primer:template hybrid used here had a C.A mismatch at the 3' terminus. V_{max} and K_m were determined by fitting the points to a “Single rectangular 2-parameter hyperbola equation” using the Sigmaplot software.

Recipes

- Extension reaction buffer (50 ml)
 - 1 M Tris HCl (pH 8) 25 ml
 - 3 M KCl 13.3 ml
 - 1 M DTT 1 ml
 - RNase free water 10.7 ml
- 2x loading dye (10 ml)

50 mM Tris HCl (pH 6.8)	500 µl
100 mM DTT	1 ml
2% SDS	2 ml
0.05% Bromophenol blue	500 µl
10% glycerol	1 ml
RNase free water	5 ml

References

1. Achuthan, V., Keith, B. J., Connolly, B. A. and DeStefano, J. J. (2014). [Human immunodeficiency virus reverse transcriptase displays dramatically higher fidelity under physiological magnesium conditions *in vitro*](#). *J Virol* 88(15): 8514-8527.
2. Hou, E. W., Prasad, R., Beard, W. A. and Wilson, S. H. (2004). [High-level expression and purification of untagged and histidine-tagged HIV-1 reverse transcriptase](#). *Protein Expr Purif* 34(1): 75-86.
3. Mendelman, L. V., Petruska, J. and Goodman, M. F. (1990). [Base mispair extension kinetics. Comparison of DNA polymerase alpha and reverse transcriptase](#). *J Biol Chem* 265(4): 2338-2346.
4. Menendez-Arias, L. (2009). [Mutation rates and intrinsic fidelity of retroviral reverse transcriptases](#). *Viruses* 1(3): 1137-1165.
5. Rezende, L. F. and Prasad, V. R. (2004). [Nucleoside-analog resistance mutations in HIV-1 reverse transcriptase and their influence on polymerase fidelity and viral mutation rates](#). *Int J Biochem Cell Biol* 36(9): 1716-1734.
6. Svarovskaia, E. S., Cheslock, S. R., Zhang, W. H., Hu, W. S. and Pathak, V. K. (2003). [Retroviral mutation rates and reverse transcriptase fidelity](#). *Front Biosci* 8: d117-134.