


***In vitro* Colorimetric Method to Measure Plant Glutamate Dehydrogenase Enzyme Activity**

Asier Sarasketa¹, Izargi Vega-Mas¹ and Daniel Marino^{1, 2*}

¹Department of Plant Biology and Ecology, University of the Basque Country (UPV/EHU), Bilbao, Spain; ²Ikerbasque, Basque Foundation for Science, Bilbao, Spain

*For correspondence: Daniel Marino: daniel.marino@ehu.es

[Abstract] Glutamate dehydrogenase (GDH) is an NAD(H) dependent enzyme that catalyzes, *in vitro*, the reversible amination of glutamate. Here we describe how to determine spectrophotometrically GDH activity monitoring NADH evolution. This protocol is described here for *Arabidopsis thaliana* (*A. thaliana*) although it is also valid for other plant species. GDH protein is a hexamer composed, in the case of *Arabidopsis*, of a combination of GDH α , GDH β and GDH γ subunits. Every combination of subunits is possible; however, it is still barely known whether different combinations affect the enzymatic properties of the hexamers. In other species, hexamers are a combination of GDH α and GDH β but it cannot be discarded the existence of other genes since for instance GDH γ subunit in *Arabidopsis* was described in Fontaine *et al.* (2012).



Materials and Reagents

1. 4-week old *Arabidopsis thaliana* leaves and roots
2. Ultrapure water
3. Liquid N₂
4. Ice
5. Flat bottom microplates (Deltalab, catalog number: 900011.1)
6. 4 mm diameter glass beads (Glaswarenfabrik Karl Hecht) (VWR International, catalog number: 201-0278)
7. 1.2 ml deep well storage plate (Thermo Fisher Scientific, catalog number: AB-0564)
8. EVA Capband for capping 8 tubes (Micronic, catalog number: MP227B1)
9. Bio-Rad Protein Assay Dye Reagent Concentrate (Bio-Rad Laboratories, AbD Serotec[®], catalog number: 500-006)
10. NADH (Sigma-Aldrich, catalog number: 43420)
11. PVPP (Sigma-Aldrich, catalog number: 77627)
12. Glycerol (EMD Millipore Corporation, catalog number: 104092)
13. Bovine Serum Albumin (BSA) (Sigma-Aldrich, catalog number: A7906)
14. Triton X-100 (Sigma-Aldrich, catalog number: T-9284)
15. HEPES (Sigma-Aldrich, catalog number: H3375)

16. KOH (EMD Millipore Corporation, catalog number: 105029)
17. MnCl₂ (Sigma-Aldrich, catalog number: M8266)
18. EDTA (Sigma-Aldrich, catalog number: ED255)
19. EGTA (Sigma-Aldrich, catalog number: E4378)
20. Dithiothreitol (DTT) (Sigma-Aldrich, catalog number: D0632)
21. PMSF (Sigma-Aldrich, catalog number: P7626)
22. Delta-aminocaproic acid (Sigma-Aldrich, catalog number: A2504)
23. Leupeptin (Sigma-Aldrich, catalog number: L2884)
24. Tricine (Sigma-Aldrich, catalog number: T0377)
25. CaCl₂ (Panreac Applichem, catalog number: 131232)
26. 2-Oxogutarate (Merck KGaA, catalog number: 5194)
27. (NH₄)₂SO₄ (Panreac Applichem, catalog number: 131140)
28. Extraction buffer (see Recipes)
29. Reaction buffer (see Recipes)
30. Bradford solution (see Recipes)

Equipment

1. TissueLyser (RETSCH, model: MM400)
2. Plate centrifuge (Sigma, model: 2-16K)
3. Absorbance microplate reader (Biotek Powerwave X 384 Microplate spectrophotometer)
4. TissueLyser Adapter Set 2 x 96 (QIAGEN, catalog number: 69984)

Procedure

A. Material harvest and homogenization

1. In a 96 deepwell (1.2 ml) plate, place 2 glass beads (4 mm diameter) into each well.
2. Weigh approximately 20 mg of fresh leaves or roots and place them in a 96-deepwell plate, half-submerged in liquid nitrogen. Store sealed with EVA Capband at -80 °C until use.
3. Grind frozen tissue two times with a TissueLyser for 60 sec at 27 Hz frequency. Plates are coupled to the TissueLyser with a TissueLyser Adapter Set 2 x 96. (Ensure sealing of the wells during homogenization).

B. Extraction

1. Add around 3-5 mg PVPP per well (this approximately corresponds to a level micro spatula).
2. Add 500 µl of cold extraction buffer to each well and rapidly mix by inverting the plate (avoid thawing the samples prior to adding the buffer).

3. Homogenize the samples again with TissueLyser two times for 60 sec at 27 Hz frequency.
4. Centrifuge the plate at 4,000 rpm and 4 °C for 30 min in a plate centrifuge.
5. Recover the supernatants (400 µl) in a new 96 deepwell plate and keep the samples on ice.

C. Determination of protein content by Bradford method

1. In a 96-well spectrophotometer plate, place 1 µl of extract per well and add ultrapure water up to 10 µl. Make a blank with 1 µl of extraction buffer.
2. For the calibration curve place BSA standards in a range of 0 to 6 µg of BSA/well using a 1 µg/µl BSA stock and complete volume up to 10 µl.
3. Add in every well 240 µl of Bradford solution and incubate for 5 min at room temperature in the dark.
4. Read absorbance at 595 nm.

D. Enzyme activity measurement

1. Place 20 µl of supernatant per well.
2. Add 280 µl of reaction buffer. This step should be performed rapidly since the reaction starts as soon as the buffer is added. Thus, if measuring 96 samples the use of a multichannel pipette is strongly advised to ensure reproducibility.
3. Immediately monitor NADH evolution kinetic at 340 nm in the spectrophotometer for 10 min at 30 °C (Figure 1). Ensure spectrophotometer is already at 30 °C before adding the reaction buffer to the sample.




Figure 1. Graph showing NADH consumption by GDH aminating activity

4. For NADH calibration curve, place 300 μ l NADH dilutions from 0-0.2 mM in a microplate and read at 340 nm. (NADH dilutions must be prepared just prior to the measurement of the calibration curve to avoid the oxidation of NADH).

Data analysis

1. To determine protein concentration in the extracts, subtract from the absorbance of every sample the absorbance of the blank (done with extraction buffer) and calculate the protein concentration with the calibration curve performed with BSA standards.
2. Enzyme activity is calculated with the maximum speed (V_{max}) obtained from NADH oxidation monitoring at 340 nm (don't take the two first minutes of the measurement into account and ensure linearity of the activity). Subtract the blank (made with extraction buffer) from every sample and calculate GDH activity with NADH calibration curve. Finally express the activity per μ mol of NADH consumed per minute and per μ g of protein (obtained from Bradford assay) or per mg of fresh/dry weight.

Notes

1. This protocol can be easily adapted to Eppendorf tubes and spectrophotometer cuvettes instead of 96-well plates just scaling the volumes. Similarly the homogenization of the samples might be done with a mortar and pestle instead of the TissueLyser.
2. If possible, we advice to do every 96-well plate step using multi-channel pipettes to save time and thus to gain reproducibility among samples.
3. This protocol is for *Arabidopsis thaliana* but it can be used for every species. For some species desalting of protein extracts supernatants might be considered.

Recipes

Note: All solutions should be made in ultrapure water.

1. Extraction buffer
 - Glycerol 10%
 - BSA 0.05%
 - Triton X-100 0.1%
 - 50 mM HEPES (pH 7.5, adjusted with KOH)
 - 10 mM MgCl₂
 - 1 mM EDTA
 - 1 mM EGTA
 - 1 mM PMSF
 - 10 mM DTT
 - 1 mM Delta-aminocaproic acid
 - 1 μM Leupeptin
2. Reaction buffer (pH 8)
 - 100 mM Tricine (pH 8, adjusted with KOH)
 - 1 mM CaCl₂
 - 50 mM (NH₄)₂SO₄
 - 0.25 mM NADH
 - 13 mM 2-Oxoglutarate
3. Bradford solution
 - Dilute Bio-Rad Protein Assay Dye Reagent Concentrate 1:5 with ultrapure water
 - Stored in a dark bottle at 4 °C

Acknowledgments

This protocol is adapted from Sarasketa *et al.* (2014) and based on the methodology reported by Groat and Vance (1981). This work was supported by the Basque Government (IT526-10), the UPV/EHU (EHUA14/14), the People Program (Marie Curie Actions) of the European Union's Seventh Framework Program (FP7/2007–2013) under REA grant agreement number 334019 and MINECO (BIO2014-56271-R).

References

1. Bradford, M. M. (1976). [A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding.](#) *Anal Biochem* 72: 248-254.
2. Fontaine, J. X., Terce-Laforgue, T., Armengaud, P., Clement, G., Renou, J. P., Pelletier, S., Catterou, M., Azzopardi, M., Gibon, Y., Lea, P. J., Hirel, B. and Dubois, F.

-
- (2012). [Characterization of a NADH-dependent glutamate dehydrogenase mutant of *Arabidopsis* demonstrates the key role of this enzyme in root carbon and nitrogen metabolism.](#) *Plant Cell* 24(10): 4044-4065.
3. Groat, R. G. and Vance, C. P. (1981). [Root nodule enzymes of ammonia assimilation in alfalfa \(*Medicago sativa* L.\): DEVELOPMENTAL PATTERNS AND RESPONSE TO APPLIED NITROGEN.](#) *Plant Physiol* 67(6): 1198-1203.
 4. Sarasketa, A., Gonzalez-Moro, M. B., Gonzalez-Murua, C. and Marino, D. (2014). [Exploring ammonium tolerance in a large panel of *Arabidopsis thaliana* natural accessions.](#) *J Exp Bot* 65(20): 6023-6033.