

Bone Marrow Mesenchymal Stem Cells Adhesion Assay

Zhigang Yang and Ran Xiao*

Research Center of Plastic Surgery Hospital, Chinese Academy of Medical Sciences & Peking Union of Medical College, Beijing, China

*For correspondence: xiaoran@psh.pumc.edu.cn

[Abstract] Mesenchymal stem cells (MSCs) are widespread in adult organisms and involved in tissue maintenance and repair as well as in the regulation of hematopoiesis and immunologic responses. As cell adhesion play important roles in cell interactions and signaling, to thoroughly evaluate the adhesion ability of MSCs is of vital important to clarify the mechanism of self-renewal, proliferation, activation and migration of MSCs in different microenvironments. Based on the method by Siler *et al.*, 2000, we revised the protocol in order to provide details on how to evaluate the adhesion ability of MSCs from bone marrow (BMSCs) on extracellular matrix (ECM) protein laminins. The current protocol can also be easily translated to MSCs with other treatments or ECMs such as collagens, fibronectin, *etc.*

Materials and Reagents

1. 96 well culture plate (Sigma-Aldrich, Corning® Costar®, catalog number: CLS3595)
2. Human bone marrow derived MSCs, isolated and cultured as previously described (Kern *et al.*, 2006)
Note: Cells in the third cell passage were used.
3. Laminin 511&521 (Biolamina, catalog number: LN511; LN521)
4. DMEM (Thermo Fisher Scientific, Invitrogen™, catalog number: 11965092)
5. Fetal bovine serum (FBS) (Thermo Fisher Scientific, Invitrogen™, catalog number: 10099141)
6. Trypsin (Thermo Fisher Scientific, Invitrogen™, catalog number: 25200-056)
7. Soybean trypsin inhibitor (Thermo Fisher Scientific, Invitrogen™, catalog number: 17075-029)
8. BSA (Sigma-Aldrich, catalog number: V900933)
9. Blocking buffer (0.5% BSA in PBS without Ca²⁺ and Mg²⁺)
10. Wash buffer (0.1% BSA in PBS without Ca²⁺ and Mg²⁺)
11. 2% SDS (Sigma-Aldrich, catalog number: 71729)
12. 4% paraformaldehyde (Sigma-Aldrich, catalog number: 158127)
13. NaCl (Sigma-Aldrich, catalog number: S7653)
14. KCl (Sigma-Aldrich, catalog number: 746436)
15. Na₂HPO₄ (Sigma-Aldrich, catalog number: 795410)
16. KH₂PO₄ (Sigma-Aldrich, catalog number: P0662)
17. CaCl₂·2H₂O (Sigma-Aldrich, catalog number: 223506)

18. MgCl₂·6H₂O (Sigma-Aldrich, catalog number: M9272)
19. HCl (Sinopharm chemical reagent Being Co.ltd, catalog number: 10011008)
20. Crystal violet powder (Sigma-Aldrich, catalog number: C6158)
21. Ethanol (Sinopharm chemical reagent Being Co.ltd, catalog number: 10009159)
22. PBS with/without Ca²⁺ and Mg²⁺ (see Recipes)
Note: Commercial DPBS with/without calcium and magnesium can also be used.
23. 0.1% crystal violet staining solution (see Recipes)

Equipment

1. Small shaker for microtiter plate (IKA, model: MS3 Digital)
2. Scanner (UMAX, model: POWERLOOK 2100XL)
3. Series II 3110 Water-Jacketed CO₂ chamber (Thermo Fisher Scientific, Forma™, catalog number: 3111)
4. Microscope (Nikon, model: ECLIPSE TE2000-S)
5. Microplate reader (ThermoFisher Scientific, model: Multiscan MK3) or spectrometer with 550 nm wavelength available.

Procedure

1. Coat plate with laminins
 - a. Slowly thaw recombinant laminins at 2 °C to 8 °C before use.
 - b. Dilute the thawed laminin stock solution with 1x DPBS (Ca²⁺/Mg²⁺) to 10 µg/ml, Add 60 µl diluted laminin solution to each well of 96 well culture plate (The final coating concentration is 2 µg/cm²). Make sure the laminin solution is spread evenly across the surface. Leave some wells uncoated as negative control.
 - c. Incubate at 2 °C to 8 °C overnight.
 - d. Wash with wash buffer (200 µl/well) for 2 times.
 - e. Block plates with blocking buffer (200 µl/well) at 37 °C in CO₂ chamber for 60 min.
 - f. Wash with wash buffer (200 µl/well) for 2 times.

Note: All the wash procedures do not need some time incubation, the followings are the same.
2. Prepare and seed cells
 - a. Taking a 10 cm culture dish as an example, wash the cells with 2 ml PBS.
 - b. Detach the cells with 2 ml 0.25% Trypsin-EDTA, neutralize trypsin with 2 ml 0.25 mg/ml soybean trypsin Inhibitor.
 - c. Mix the cells by pipetting up and down using a 1 ml pipette, then transfer them to a 15 ml sterile tube and centrifuge at 300 x g for 5 min.
 - d. Resuspend the cells in 2 ml DMEM with 0.5% FBS.

- e. Count the cells and dilute to a final concentration of 2×10^5 /ml, add 100 μ l (2×10^4 cells) to each well. Set up at least triplicate wells for each condition.

Note: Cell number vary depending on the coated ECM, for laminins, 2×10^4 is appropriate. However, 4×10^5 cells may be required for non-coated wells.
 - f. Incubate at 37 °C (in a CO₂ chamber) for 20 min.

Note: Adhesion time depends on the coated ECM. It is useful to observe the adherent condition under microscope at different time points, choosing between 5 min and 30 min (5, 10, 15, 20, 30), even do a time course first. In our system, cells adhere to the laminin-coated wells within at least 10 min, while for non-coated wells, it may take longer time.
3. Stop the assay
 - a. Discard the seeding medium carefully without scratching the bottom of wells.
 - b. Rinse cells once with 200 μ l PBS per well.
 - c. Shake the plate at 2,000 rpm for 15 sec.
 - d. Discard the PBS and wash with 200 μ l wash buffer per well for 3 times.
 - e. Fix with 200 μ l 4% paraformaldehyde per well. Incubate at room temperature for 10-15 min.
 - f. Wash with 200 μ l PBS per well.
 4. Staining and analysis
 - a. Add 100 μ l 0.1% crystal violet to each well and incubate for 10 min at room temperature.
 - b. Remove the crystal violet and wash with ddH₂O (200 μ l/well) for 3 times.
 - c. Turn the plates upside down on the absorbent papers. Let the plates dry up completely.
 - d. The plates can be scanned by a scanner firstly.
 - e. Add 100 μ l 2% SDS to each well and incubate for 10 min with gently shaking at room temperature.
 - f. Read the absorbance within 5 min on a microplate reader/spectrometer at a wavelength of 550 nm.

Representative data




Figure 1. Laminin 511 promotes the adhesion of bone marrow mesenchymal stem cells (BMSCs). A. The scanned image of crystal violet staining of BMSCs adhered to laminin 511 and BSA coated wells; B. The O.D. values from the dissolved crystals are shown for (A), the data are presented as the mean \pm SEM from three replicate wells; ***, $P < 0.001$.

Recipes

1. PBS with/without Ca²⁺ and Mg²⁺ (Reference 1)
 - a. Dissolve the following in 800 ml distilled H₂O
 - 8 g NaCl
 - 0.2 g KCl
 - 1.44 g Na₂HPO₄
 - 0.24 g KH₂PO₄For PBS with Ca²⁺ and Mg²⁺, supplement with the following
 - 0.133 g CaCl₂·2H₂O
 - 0.10 g MgCl₂·6H₂O
 - b. Adjust pH to 7.4 with HCl
 - c. Adjust volume to 1 L with additional distilled H₂O
 - d. Sterilize by autoclaving
2. 0.1% crystal violet staining solution
To make up 50 ml crystal violet solution, dissolve 50 mg crystal violet powder in (5 ml ethanol/45 ml water).

References

1. [Phosphate-buffered saline \(PBS\)](#). (2006) *Cold Spring Harb Protoc*: doi:10.1101/pdb.rec8247.
2. Kern, S., Eichler, H., Stoeve, J., Kluter, H. and Bieback, K. (2006). [Comparative analysis of mesenchymal stem cells from bone marrow, umbilical cord blood, or adipose tissue](#). *Stem Cells* 24(5): 1294-1301.
3. Siler, U., Seiffert, M., Puch, S., Richards, A., Torok-Storb, B., Muller, C. A., Sorokin, L. and Klein, G. (2000). [Characterization and functional analysis of laminin isoforms in human bone marrow](#). *Blood* 96(13): 4194-4203.