

***Schistosoma haematobium* Egg Isolation**

Chi-Ling Fu and Michael H. Hsieh*

Urology Department, Stanford University, Stanford, USA

*For correspondence: mhsieh@childrensnational.org

[Abstract] *Schistosoma haematobium* is the etiologic agent for urogenital schistosomiasis, a major source of morbidity and mortality for more than 112 million people worldwide. Infection with *S. haematobium* results in a variety of immunopathologic sequelae caused by parasite oviposition within the urinary tract, which drives inflammation, hematuria, fibrosis, bladder dysfunction, and increased susceptibility to urothelial carcinoma. Since most of the pathology in schistosomiasis is directly attributable to the host reaction to eggs and egg-associated antigens, their isolation and study are important experimental techniques. *S. haematobium* eggs can be collected from infected tissues for injection into other animals or preparation of crude egg extracts. This protocol describes a simple way to isolate eggs.

Schistosomes are a biohazard. Workers should wear latex gloves at all times when handling schistosomal materials or any tissues from infected animals.

Materials and Reagents

1. *Schistosoma haematobium*-infected hamster tissues (intestines and livers)
2. NaCl
3. 1.2% (w/v) NaCl (autoclaved) (see Recipes)
4. 0.85% (w/v) NaCl (autoclaved) (see Recipes)

Equipment

1. Waring blender, with variable speed control (Eberbach, model: 8580 semi-micro container and 2-speed Waring power base unit)
2. Stainless steel sieves (Newark Wire Cloth sieve No. 40, 80, 140, 325; mesh openings of 420 μm , 180 μm , 105 μm , and 45 μm)
3. Plastic spray bottle (e.g. Curtin-Matheson)
4. Glass Petri dishes with flat bottoms (Corning Incorporated, 100 x 20 mm)
5. Pasteur pipets
6. Hematocytometer
7. Microscope

Procedure

1. Sacrifice hamsters at the point of maximal liver and intestinal *Schistosoma* egg levels (18 weeks post-infection).
2. Mince the infected tissues (intestines and livers) with scissors, suspend in 1.2% NaCl, and then place in a Waring blender. Blend tissues for 30 sec at low speed. Sieves are assembled from top to bottom (420 μm , 180 μm , 105 μm , and 45 μm) (see Figure 1) with progressively decreasing pore sizes. Pass the blended material through the crudest sieve (420 μm pores).
3. Collect the tissue remaining on top of the sieve and blend again for 30 sec, this time at medium speed.
4. Pass the resulting material through the sieves as before, collect the remaining tissue from the upper surface of the sieve, and blend for 30 sec at the highest speed setting. Pass the tissue through the crude sieve an additional time.
5. Pass the filtrate through the series of stainless steel sieves with decreasing pore sizes, using a plastic spray bottle filled with 1.2% NaCl to rinse the filtrate through the successive screens.
6. Following repeated washes through the upper sieves, the eggs will be retained on the final sieve (pore size of 45 μm).
7. In order to reduce contamination with residual hamster tissues, prepare a second set of sieves and pass the egg-containing solution through the crude sieve (420 μm opening) and repeat steps 4 and 5.
8. Wash the eggs off the final sieve into a glass Petri dish. With gentle swirling, concentrate the eggs in the middle of the dish (eggs look like grains of brown sand by the naked eye), aspirate with a Pasteur pipet, and place them in a tube on ice. Add enough 1.2% NaCl to the Petri dish each time when eggs are removed to keep the volume constant, and repeat swirling and concentrating the eggs until no more can be concentrated in the middle of the dish.
Note: For best results eggs should be collected using a Petri dish with a perfectly flat bottom.
9. When no more eggs can be concentrated in the middle of the Petri dish upon swirling, discard the contents of the dish, place the collected eggs back into the dish with fresh 1.2% NaCl and repeat the swirling technique, and collect the concentrated eggs as before.
10. Finally, count eggs using a hemacytometer.

After several series of these purification steps, most of the tissue debris and immature eggs will be eliminated. The final egg population will be enriched for mature eggs.

Although it is best to use fresh eggs, eggs can be kept at 4 °C for several days in 0.85% NaCl without appreciable loss of viability.



Figure 1.

Recipes

1. 1.2% (w/v) NaCl (autoclaved)
2. 0.85% (w/v) NaCl (autoclaved)

Note: For downstream in vivo studies, Antibiotic-Antimycotic solution (Life Technologies, Gibco®) can be added to the saline solution used in the final step.

Acknowledgments

We acknowledge the following citations from which this protocol was adapted: Fu *et al.* (2011); Fu *et al.* (2012). We also acknowledge NIH DK087895 for their funding support.

References

1. Fu, C. L., Apelo, C. A., Torres, B., Thai, K. H. and Hsieh, M. H. (2011). [Mouse bladder wall injection](#). *J Vis Exp* (53): e2523.
2. Fu, C. L., Odegaard, J. I., Herbert, D. R. and Hsieh, M. H. (2012). [A novel mouse model of Schistosoma haematobium egg-induced immunopathology](#). *PLoS Pathog* 8(3): e1002605.
3. Lewis, F., Schistosomiasis, in Current Protocols in Immunology. 2001, John Wiley & Sons, Inc.