

## ***In vitro* RNA-protein Binding Assay by UV Crosslinking**

Hailong Zhang\* and Muxiang Zhou

Department of Pediatrics, Emory University, Atlanta, USA

\*For correspondence: [hzhan52@emory.edu](mailto:hzhan52@emory.edu)

**[Abstract]** Because covalent bond can form between RNA and its binding proteins after UV irradiation, UV cross-linking is widely used to identify the specific RNA binding proteins. This protocol is described in details as follows.

### **Materials and Reagents**

1. Linearized DNA template for transcription of the RNA of interest
2. MAXIscript *in vitro* Transcription Kit (Applied Biosciences, catalog number: AM1308M)  
(This kit contain ATP, CTP, GTP,  $\alpha$ -P32UTP and DNase I)
3. RNasin (Promega Corporation, catalog number: N2611)
4. RNase T1 (Applied Biosciences, catalog number: 10109193001)
5. 0.5 M EDTA (Life Technologies, Gibco<sup>®</sup>, catalog number: 15575-038)
6. ddH<sub>2</sub>O (GeneMate Loyalty, catalog number: UPW-1000)
7. HEPES (Sigma-Aldrich, catalog number: H3375)
8. MgCl<sub>2</sub> (Sigma-Aldrich, catalog number: M-2393)
9. Glycerol (Sigma-Aldrich, catalog number: G7893-1L)
10. DTT (Sigma-Aldrich, catalog number: D9779-5G)
11. Tris-Base (Thermo Fisher Scientific, catalog number: BP152-1)
12. SDS (Sigma-Aldrich, catalog number: L-4390)
13. Bromphenol blue (Sigma-Aldrich, catalog number: B5525)
14. TEMED
15. Ammoniampersulfate
16. 5x binding buffer (see Recipes)
17. 2x SDS loading buffer (see Recipes)
18. 10x TGE buffer (see Recipes)
19. 6% TGE gel (see Recipes)

### **Equipment**

1. SPIN-Pure<sup>TM</sup> column (G-50) (PireBiotech SCW50-50 DEPC-water)

2. SDS-PAGE system (Bio-Rad, catalog number: 165-1802)
3. UV Stratalinker 1800 (Stratagene, catalog number: 474645)
4. Film processor (Konica Minolta, model: SRX-101A)
5. Centrifuges (Eppendorf, catalog number: 5810R)
6. Scintillation Counter (Beckman, catalog number: LS6500)
7. Bench top Radiation Shield with 6-1/4" base (Cole-Parmer, catalog number: WU-36218-00)
8. 37 °C water bath

## **Procedure**

- A. *In vitro* transcription of <sup>32</sup>P-labeled RNA probes
  1. DNA templates for synthesis of the RNA probe are generated by PCR using specific primers.
  2. Thaw frozen reagents, place RNA polymerase on ice and vortex the 10x Transcription buffer and keep the buffer at room temperature. All reagents should be centrifuged briefly before opening.
  3. Assemble transcription reaction at room temperature as the following sequence:
    - 1 µg DNA template
    - ddH<sub>2</sub>O to 20 µl
    - 2 µl 10x transcription buffer (after addition of water and DNA)
    - 1 µl 10 mM ATP
    - 1 µl 10 mM CTP
    - 1 µl 10 mM GTP
    - 5 µl [α-<sup>32</sup>P]UTP
    - 2 µl T7 enzyme mix
    - 2 µl RNasin (40 U/µl)
 Mix thoroughly.
  4. Incubate 1 h at 37 °C.
  5. Add 1 µl of 5 mg/ml DNase I to digest the rest template DNA and incubate 10 min in 37 °C water bath.
  6. Add 1 µl of 0.5 M EDTA, mix and add 78 µl ddH<sub>2</sub>O to adjust final volume to 100 µl and mix.
- B. Purification of <sup>32</sup>P-labeled RNA probes with SPIN-Pure column (G-50)
  1. Allow a minimum of 30 min of room temperature to warm the columns before the following steps.
  2. Gently invert the column several times to suspend the column buffer (DEPC-Water).

3. Remove the top cap from the column, and then remove the bottom tip.
4. Allow the column buffer to drain by gravity before proceeding.
5. Place this column/tube apparatus into an adaptor tube. Centrifuge at 1,100 x g for 2 min at room temperature.
6. Repeat centrifugation again to let the column dry completely and discard the collection tube and the eluted buffer.
7. Put the column in a second collection tube in upright position and apply the RNA sample (20 to 50  $\mu$ l) to the center of the column gel very slowly and carefully (the rest RNA sample can be stored at -80 °C for one week).
8. Centrifuge at 1,100 x g for 4 min. The purified  $^{32}$ P-labeled RNA probe is collected in the bottom of the collection tube. Discard the spin column and continue with the following procedure.
9. Pick 1  $\mu$ l of the RNA sample and count the cpm value.


#### C. RNA and protein binding reaction

1. Prepare the following mixture on ice:
  - $^{32}$ P-labeled RNA probe ( $2 \times 10^5$  cpm)
  - 1  $\mu$ l 12.5 mM ATP
  - 3  $\mu$ l 5x binding buffer
  - 3  $\mu$ l 0.5 M KCl
  - Protein (100 ng) (this protein can be a GST fusion protein purified according to the protocol prepared by GE Healthcare Life Sciences)
  - H<sub>2</sub>O to final volume of 15  $\mu$ l.
2. Mix gently by pipetting up and down for 10 times, avoiding vortex.
3. Place the uncovered tube containing the reaction mixture on ice, directly underneath the bulb (about 10 cm from the surface) of a 254-nm UV light source, irradiation with  $4 \times 10^5$   $\mu$ J/cm<sup>2</sup> energy.
4. Incubate at room temperature for 30 min then keep on ice for another 3 h.

#### D. Remove the non-incorporated $^{32}$ P

1. Add 1  $\mu$ l of RNase T1 (1 U/ $\mu$ l) to each tube and incubate for additional 10 min at 37 °C to degrade the free RNA.
2. Add an equal volume (16  $\mu$ l) of 2x SDS loading dye and separate the sample in TGE-gel or SDS-PAGE (no boiling need).
3. Autoradiography for 12 h to 7 days for visualization.

Principle Figure



## Recipes

1. 5x binding buffer
  - 50 mM HEPES (pH 7.2)
  - 15 mM MgCl<sub>2</sub>
  - 25% glycerol
  - 5 mM DTT
  - Stored at -20 °C
2. 2x SDS loading buffer
  - 125 mM Tris-Cl (pH 7.6)
  - 20% (v/v) glycerol
  - 4% (w/v) SDS
  - 0.008% (w/v) bromphenol blue
  - 20 mM DTT
3. 10x TGE buffer
  - Tris 100 mM
  - Glycine 1 M
  - EDTA 10 mM
  - pH 8.3
4. 6% TGE gel
  - 6 ml 30% acrylamide
  - 5 ml 10x TGE buffer
  - 35 ml H<sub>2</sub>O
  - 300 µl ammoniampersulfate
  - 22 µl TEMED

### Notes:

- a. *Avoiding RNase contamination is the key for successful.*

- 
- b. As  $^{32}\text{P}$  is radioactive, this experiment should be performed under the protection of Benchtop Radiation Shield.

### **Acknowledgments**

This work was supported by the National Institutes of Health (R01 CA123490 and R01CA143107 to MZ) and CURE (MZ and LG).

### **References**

1. Gu, L., Zhang, H., He, J., Li, J., Huang, M. and Zhou, M. (2012). [MDM2 regulates MYCN mRNA stabilization and translation in human neuroblastoma cells](#). *Oncogene* 31(11): 1342-1353.
2. Walker, J., de Melo Neto, O. and Standart, N. (1998). [Gel retardation and UV-crosslinking assays to detect specific RNA-protein interactions in the 5' or 3' UTRs of translationally regulated mRNAs](#). *Methods Mol Biol* 77: 365-378.