

Site-directed Mutagenesis Using Dpn1

Lili Jing*

Department of Cell and Molecular Biology, University of Pennsylvania, Philadelphia, USA

*For correspondence: lilijingcn@gmail.com

[Abstract] Site-directed mutagenesis is an important and widely used tool in molecular biology to generate specific changes in the DNA sequence of a given gene/genome. The protocol described here is for making nucleotide changes at specific loci in a large vector (≥ 10 kb), and is based on the QuikChange II XL Site-Directed Mutagenesis Kit.

Materials and Reagents

1. Dpn1 restriction enzyme
2. XL10-Gold Ultracompetent Cells
3. QuikChange II XL Site-Directed Mutagenesis Kit (Guidechem)
4. LB agar (Sigma-Aldrich)
5. Antibiotics (Sigma-Aldrich)
6. NZ amine (casein hydrolysate)
7. Yeast extract
8. NaCl
9. NaOH
10. MgCl₂
11. MgSO₄
12. Glucose
13. NZY+ broth (see Recipes)

Equipment

1. Thermal cyclers (Bio-Rad Laboratories)
2. Bench-top centrifuge

Procedure

A. Mutant strand synthesis reaction

1. Prepare the sample reaction as follows:

- 5 μ l of 10x reaction buffer
- X μ l (10 ng) of dsDNA template
- 1.25 μ l (125 ng) of oligonucleotide primer 1
- 1.25 μ l (125 ng) of oligonucleotide primer 2
- 1 μ l of dNTP mix
- 3 μ l of QuikSolution
- ddH₂O to a final volume of 50 μ l
- Then add 1 μ l of PfuUltra HF DNA polymerase (2.5 U/ μ l)

2. Cycle each reaction using the following parameters (be sure to adhere to the 18 cycle limit):

Segment	Cycles	Temperature	Time
1	1	95 °C	1 min
2	18	95 °C	50 sec
		60 °C	50 sec
		68 °C	1 min/kb of plasmid length*
3	1	68 °C	7 min

*For example, a 5 kb plasmid requires 5 min at 68 °C per cycle.

3. Following cycling, place reaction tubes on ice for 2 min to cool the reactions to 37 °C.
4. Check product by electrophoresis of 10 μ l of product on 1% agarose gel. While gel is running, start Dpn1 digestion.

B. Dpn1 Digestion of the Amplification Products

1. Add 1 μ l of the Dpn1 restriction enzyme (10 U/ μ l) directly to each amplification reaction using a small, pointed pipet tip.
2. Gently mix each reaction mixture by pipetting up and down several times. Spin for 1 minute and incubate the reaction at 37 °C for 1 h to digest the parental supercoiled dsDNA.

C. Transformation of XL10-Gold Ultracompetent Cells.

1. Gently thaw XL10-Gold ultracompetent cells on ice. For each control and sample reaction to be transformed, aliquot 45 μ l of the cells to a pre-chilled polypropylene tube.
2. Add 2 μ l of the B-ME mix provided with the kit to 45 μ l cells.
3. Swirl the contents of the tube gently. Incubate on ice for 10 min, swirling gently every 2 min.
4. Transfer 2 μ l of the Dpn1-treated DNA from each control and sample reaction to separate aliquots of the ultracompetent cells.

- For a positive control, add 1 μ l of 0.01 ng/ μ l pUC18 control plasmid.
5. Preheat NZY+ broth in a 42 °C water bath.
 6. Heat-pulse tubes in a 42 °C water bath for 30 sec.
 7. Incubate the tubes on ice for 2 min.
 8. Add 0.5 ml of preheated (42 °C) NZY+ broth to each tube, then incubate the tubes at 37 °C for 1 h with shaking 225-250 rpm.
 9. Plate the appropriate volume of each transformation reaction as indicated in the table below on the appropriate selection plates.

Reaction type	Volume to plate
pWhitescript mutagenesis control	250 μ l
pUC18 transformation control	5 μ l (in 200 μ l NZY+ broth)
Sample mutagenesis	250 μ l on each of two plates (entire transformation reaction)

10. Incubate plates at 37 °C for >16 h.

Recipes

1. NZY+ Broth (per Liter)
 - 10 g of NZ amine (casein hydrolysate)
 - 5 g of yeast extract
 - 5 g of NaCl
 - Add deionized H₂O to a final volume of 1 L
 - Adjust to pH 7.5 using NaOH and then autoclave
 - Add the following filter-sterilized supplements prior to use:
 - 12.5 ml of 1 M MgCl₂
 - 12.5 ml of 1 M MgSO₄
 - 20 ml of 20% (w/v) glucose (or 10 ml of 2 M glucose)

Acknowledgments

This protocol was developed in the Michael Granato Lab at University of Pennsylvania, Philadelphia, USA, and modified from the manual of QuikChange II XL Site-Directed Mutagenesis Kit. This work was supported by NIH grant R01HD037975.