

## DAB Staining and Visualization of Hydrogen Peroxide in Wheat Leaves

Zhaohui Liu\* and Tim Friesen

Department of Plant Pathology, North Dakota State University, Fargo, USA

\*For correspondence: [zhliu@ndsu.edu](mailto:zhliu@ndsu.edu)

**[Abstract]** The production of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) has been recognized as an important feature of plant cells that undergo programmed cell death (PCD) during host-pathogen interaction. Thordal-Christensen *et al.* (1997) first described a method using chemical 3,3-diaminobenzidine (DAB) to detect the presence and distribution of H<sub>2</sub>O<sub>2</sub> in barley leaves challenged by the *powdery mildew fungus* (Thordal-Christensen *et al.*, 1997). Since then, this method has been adapted to many other plant species for *in situ* detection of H<sub>2</sub>O<sub>2</sub>. Here, we describe a modified protocol to stain and visualize H<sub>2</sub>O<sub>2</sub> production in wheat leaves during infection by the necrotrophic fungus, *Stagonospora nodrum* or infiltration by the necrotrophic effectors produced by the fungus. The short version of this method has been reported in Liu *et al.* (2012).

### Materials and Reagents

1. 3,3-diaminobenzidine (Sigma-Aldrich, catalog number: D8001)
2. 0.1 N HCl
3. 95% ethanol (bulk)
4. Glacial acetic acid (EMD Millipore, catalog number: UN2789)
5. L-(+)-Lactic acid (Sigma-Aldrich, catalog number: L1750)
6. Glycerol (Life Technologies, Invitrogen™, catalog number: 15514-011)
7. Aluminum foil
8. Round plastic screens with the diameter closed to that of 15 ml tube (handmade, help the wheat leaf segments immerse in the staining solution). These plastic screens were cut from a big piece of plastic mesh that can be purchased from a local hardware store or online store (for example, Industrial Netting)
9. Wheat leaves at three leaf stage (~2 weeks old grown under the greenhouse conditions)

### Equipment

1. Polyethylene 50 ml conical centrifuge tube (BD Biosciences, Falcon®, catalog number: C2745)

2. Polyethylene 15 ml conical centrifuge tube (BD Biosciences, Falcon<sup>®</sup>, catalog number: C2750)
3. Glass petri dish 100 mm with lid (Sigma-Aldrich, catalog number: CLS3160102)
4. Balance
5. pH meter
6. Shaker
7. Vacuum
8. Microscope
9. Dessicator

### **Procedure**

#### A. Preparation of fresh DAB staining solution

1. Measure ~50 ml distilled water and add 0.1 N HCl (1-2 drops) till pH reaches 3.6.
2. Pour 10 ml of the above solution into each 15 ml tube (the number of tubes needed is dependent on the number of treatments).
3. Weigh 10 mg 3, 3-diaminobenzidine and add to each tube.
4. Close the cap, wrap the whole tube with Aluminum foil, and vigorously shake at 37 °C in an orbital shaker for at least 1 h (leaf samples can be collected during this time period).
5. Check the solution to see if most of the DAB is dissolved. The solution should be colorless or slight pink and should be used fresh for staining.

#### B. Staining of wheat leaves

1. Collect wheat leaves at the designated time points after inoculation with the pathogen or infiltration with toxins. The fungal inoculation and toxin infiltration in this case were done at three leaf stage.
2. Cut infiltrated or inoculated areas of the leaves into 3 cm long segments and immerse them in the DAB solution with a plastic screen (no more than 8 leaf segments in one 15 ml tube).
3. Place all tubes with the caps removed in a vacuum dessicator and vacuum for 30 min at room temperature. It's not necessary to keep the tubes in the dark at this step.
4. Release vacuum and place all tubes in a dark cabinet with the caps closed and incubate overnight.
5. The necrotic areas caused by the fungus or the toxin should be dark brown in color.

#### C. Clearing of leaves for examination (all procedures should be carried out in a fume hood)

1. Gently rinse DAB solution off the stained leaf segments using distilled water and dry slightly on paper towels.
2. Place the leaf segments on paper towels saturated with a fixative solution (ethanol/acetic acid 3:1, V/V) in a petri dish. A little amount of extra solution left in petri dish is allowed.
3. Incubate the leaf segments at room temperature in a fume hood for at least 24 h (it may require additional time to clear all the chlorophyll from the leaf tissue).
4. Transfer the cleared leaf segments to paper towels saturated with water and leave for 30 min to remove any fixative residues.
5. Transfer leaf segments on paper towels saturated with a lactoglycerol solution (lactic acid/glycerol/H<sub>2</sub>O 1:1:1, v/v/v). A little amount of extra solution left in petri dish is allowed.
6. Leaf segments can be subjected to examination after a couple of hour incubation at room temperature or stored on paper towel saturated with the same lactoglycerol solution for several weeks.
7. For examination, carefully mount leaf segments on a glass slide and observe under the microscope to determine the cellular location of DAB. Photographs can be made using a white background.

### **Acknowledgments**

This protocol was adapted from a method published in Liu *et al.* (2012), which was originally based on the descriptions by Thordal-Christensen *et al.* (1997). Development and implementation of this protocol was funded by USDA-NIFA AFRI Microbial Biology Program - Competitive grant #2010-65108-20543, and by USDA-ARS CRIS Project #5442-22000-048-0D.

### **References**

1. Liu, Z., Z. Zhang, J. D. Faris, R. P. Oliver, R. Syme, M. C. McDonald, B. A. McDonald, P. S. Solomon, S. Lu, W. L. Shelver, S. Xu and T. L. Friesen (2012). [The cysteine rich necrotrophic effector SnTox1 produced by \*Stagonospora nodorum\* triggers susceptibility of wheat lines harboring Snn1](#). *PLoS Pathog* 8(1): e1002467.
2. Thordal-Christensen, H., Z. Zhang, Y. Wei and D. B. Collinge (1997). [Subcellular localization of H<sub>2</sub>O<sub>2</sub> in plants. H<sub>2</sub>O<sub>2</sub> accumulation in papillae and hypersensitive response during the barley—powdery mildew interaction](#). *Plant J* 11(6): 1187-1194.