

## Immunoelectron Microscopy in *Chlamydomonas* Cells

Maria A. Sinetova\* and Alexandra G. Markelova

Group of Ecophysiology of Microalgae, Institute of Plant Physiology RAS, Moscow, Russia

\*For correspondence: [maria.sinetova@mail.ru](mailto:maria.sinetova@mail.ru)

**[Abstract]** The method of immunoelectron microscopy is intended for localization of proteins inside the cells of *Chlamydomonas reinhardtii* or other microalgae and cyanobacteria. This protocol was used to study localization of carbonic anhydrase Cah3 with antibodies raised in rabbit, though it can be used to localize any other abundant protein. Primary rabbit antibodies are recommended because they react quickly and specifically with proteins of *C. reinhardtii*. If primary antibodies other than rabbit are used, the blocking procedure and time of incubation with primary and secondary antibodies should be adjusted.

### **Materials and Reagents**

1. Culture of *C. reinhardtii* 137mt+ (WT) (we used the strain IPPAS D-298 from the Collection of microalgae of the Institute of Plant Physiology RAS, Moscow).
2. Primary antibodies raised in rabbits against the protein of interest (we used antibodies raised against the recombinant Cah-3 protein ( $\alpha$ -CA) of *C. reinhardtii*, Agrisera, Vännäs, Sweden)
3. Secondary antibodies: Anti-Rabbit IgG conjugated with 10 nm colloidal gold particles (Sigma-Aldrich, catalog number: G7402)
4. Paraformaldehyde (Sigma-Aldrich, catalog number: P6148)
5. LR White embedding kit (Sigma-Aldrich, catalog number: 62662-1EA-F)
6. Gelatin capsules (SPI supplies, catalog number: 02308-SS)
7. Formvar and carbon-coated nickel grids (SPI supplies, catalog number: 3430N-CF)
8. Bovine serum albumin (BSA) (Sigma-Aldrich, catalog number: A7906)
9. Goat serum (Sigma-Aldrich, catalog number: G9023)
10. Rabbit serum (Sigma-Aldrich, catalog number: R4505)
11. Paraformaldehyde (Sigma-Aldrich, catalog number: P6148)
12. Ethanol
13. Uranyl acetate (e.g. SPI supplies, catalog number: 02624-AB)
14. Lead citrate (e.g. Sigma-Aldrich, catalog number: 15326)
15. NaOH
16. Phosphate buffer saline (PBS) (pH 7.4) (see Recipes)

17. Fixation solution (4% paraformaldehyde) (see Recipes)
18. Tris-buffer saline (TBS) (pH 7.4) (see Recipes)
19. 1% BSA-TBS (see Recipes)
20. Uranyl acetate (see Recipes)
21. Lead citrate (see Recipes)

### **Equipment**

1. 96 well immunology plate (e.g. Greiner Bio-One, catalog number: 650001)
2. Pasteur pipettes with thin tips (ROTHE 306.1)
3. Thermostat unit with, at least, 37-55 °C temperature range (e.g. BINDER B28)
4. Ultramicrotome (e.g. Reichert, OMU-3, Austria)
5. Grid storage box (Sigma-Aldrich, catalog number: G6276)
6. Transmission electron microscope (e.g. Libra-120, Carl Zeiss, Germany)
7. Plastic Petri dish or Teflon plate with hydrophobic surface
8. Wet chamber

### **Procedure**

1. Sample preparation:  
Spin down the samples of *C. reinhardtii* culture in 2 ml Eppendorf tubes at 13,000 x g at room temperature for 5 min. The optimal volume of the obtained pellet, which would be sufficient for embedding in 2-3 gelatin capsules, is about 50-75 µl.
2. Fixation:  
Remove supernatant and add 0.5 ml of fixation solution (4% paraformaldehyde). For proper fixation, the minimum incubation time is 3 h at 4 °C.  
*Note: Duration of fixation can be prolonged up to several days. This fixation solution preserves membranes and most proteins, but it may dislocate starch and proteins after several weeks of storage.*
3. Embedding (all steps are carried out at room temperature):
  - a. Wash the pellet with PBS buffer in the same Eppendorf tube:  
Remove supernatant with pipette.  
Add 1 ml of PBS buffer and vortex it gently.  
Incubate for 5 min.  
Spin down as described in step 1.  
Repeat this washing step 3 times.
  - b. Dehydrate pellet with series of ethanol dilution, twice in each concentration of ethanol:

20% (v/v) 15 min+15 min

40% (v/v) 15 min+15 min

50% (v/v) 20 min+20 min

70% (v/v) 20 min+20 min

For each dehydration step, discard previous solution and add 0.5 ml of the next sequential solution with clean pipette. Incubate for the indicated time. We recommend using freshly prepared solutions pre-chilled at 4 °C. Take care not to touch or disturb the pellet during these steps.

*Note: It is handy to use a disposable plastic Pasteur pipette with a thin tip to discard the solutions. If the pellet is disturbed, centrifuge the Eppendorf tube as described in step 1. If it is necessary to interrupt the procedure at this step, the samples can be stored in 70% ethanol overnight at 4 °C.*

- c. Infiltration: Sequentially replace 70% ethanol with the mixture of 70% ethanol and LR White resin:

3:1 70% ethanol: LR White for 1 h

1:1 70% ethanol: LR White for 1 h

1:3 70% ethanol: LR White for 1 h

Next steps require the complete replacement of ethanol with LR White resin:

Replace the 1:3 ethanol/resin mixture with LR White resin alone for 1 h.

Replace it again with fresh portion of the LR White resin and leave overnight.

Next day replace it with new portion of the LR White resin and keep it for embedding.

*Note: LR White resin should be mixed with catalyst according to manufacturer's protocol for 24 h before use. With time, a resin-ethanol mixture may stratify into layers. If this happens, mix the layers gently several times with a pipette (care should be taken not to damage the pellet).*

- d. Embedding:

Put gelatin capsules into wells of 96-wells immunology plate.

Add one drop of the resin into each gelatin capsule.

Put a small piece of the pellet into the capsule – one pellet usually goes into 2-3 capsules.

Fill up capsules with resin. Avoid overflow. Close capsules carefully with their caps.

Incubate at 37 °C for 1-4 days (or at 45 °C for one day) and then at 55 °C until polymerization is completed (usually 1-3 days).


*Note: Longer incubation at lower temperature helps to receive softer block, shorter incubation at higher temperature helps to receive firmer block, but more fragile. Completeness of polymerization can be checked with a needle: The resin should be solid but not fragile.*

- e. Cut the sample into sections with a microtome and place them onto Formvar and carbon-coated nickel grids.  
*Note: Formvar+carbon coating prevents unspecific labeling and improves the quality of sections (without carbon layer sections often get wrinkled).*
4. Immunocytochemistry  
Carry out all of the following manipulations with grids on the surface of 50-100  $\mu$ l drops on a hydrophobic surface (e.g. plastic Petri dish or Teflon plate).  
*A tip: For better interaction of grid surfaces with solution, put the hydrophobic surface with drops and grids on a table that is slightly trembling due to vibration of a vortex. Adjust the intensity of the vibrations so that trembling of the drop surfaces would be noticeable.*  
*When overnight incubation of the grids is required, put them into a wet chamber at 4 °C. Do not allow grids to dry out during next steps.*
  - a. Block nonspecific interaction of antibodies on a drop of 20% goat serum in BSA-TBS for 1 h.  
*Note: If nonspecific labeling still happens, increase the proportion of the goat serum up to 50 % and BSA up to 5%.*
  - b. Put the grids onto the 100  $\mu$ l drops with properly diluted primary antibodies raised against the protein of interest. The dilution can be determined only empirically; usually it is in range of 1:16 - 1:128.  
For negative control, apply rabbit serum or, better, the pre-immune serum instead of primary antibodies to the target protein.  
Add purified protein of interest to solution of primary antibodies as positive control.  
Leave grids overnight at 4 °C. You may vary time and temperature to increase or decrease the efficiency of binding and the intensity of labeling.
  - c. Wash grids on drops with TBS 5 times, leaving them on drops for 5-10 min between changes of TBS solution.
  - d. Transfer grids onto 100  $\mu$ l drops of properly diluted (according to the protocol of manufacturer) Anti-Rabbit IgG conjugated with 10 nm colloidal gold particles in BSA-TBS and expose them for 1 h at room temperature.
  - e. Wash grids on drops with TBS 5 times, then with deionized water. These washing steps should be done by quick repeated submersion of grids into 3-5 ml of TBS and water during 1 min.
  - f. Dry grids on filter paper and put them into grid storage box.
  - g. Optional – sequentially contrast the grids with uranyl acetate and lead citrate. Prepare wet chamber with NaOH crystals and filter paper wetted with 0.2 M NaOH (Figure 1). Put grids onto 100  $\mu$ l drops of uranyl acetate for 45-60 min. Pour 50  $\mu$ l drops of lead citrate onto the same plate. Wash grids by quick repeated submersion

first into 3-5 ml of 0.2 M NaOH (5-10 times), and then – into 3-5 ml of boiled deionized water (for 1 min). NaOH solution and boiled water are needed to avoid the reaction of CO<sub>2</sub> with lead, which results in formation of insoluble crystals. Put grids onto prepared lead citrate drops for 5-7 min. Wash in the same way as performed after uranyl acetate. Dry grids on filter paper and put them into grid storage box.

5. Analysis of sections with electron transmission microscope.

*Note: Only a few gold particles are allowed to be as a background in negative and positive controls.*



**Figure 1. Wet chamber with NaOH crystals for the contrast staining**

**Recipes**

1. Phosphate buffer saline (PBS) (pH 7.4)
  - a. Prepare stock solutions:
    - Stock 1: NaH<sub>2</sub>PO<sub>4</sub> 2.76 g per 100 ml
    - Stock 2: Na<sub>2</sub>HPO<sub>4</sub> 14.2 g per 500 ml
  - b. For 0.5 L of PBS buffer mix 95 ml of the Stock 1 and 405 ml of the stock 2 and add 4.25 g of NaCl (final concentration is 0.85%)  
pH of this solution should be 7.4 (may be adjusted with NaOH)

2. Fixation solution (4% paraformaldehyde)
  - a. Heat 50 ml of PBS (pH 7.4) to 70 °C
  - b. Dissolve 2 g of paraformaldehyde (Sigma-Aldrich, P6148) with stirring in hot PBS.
  - c. Cool the solution at room temperature and filter through a filter paper  
(Keep the solution at 4 °C in darkness, use within 1 week)
3. Tris-buffer saline (TBS) (pH 7.4)
 

Tris-HCl 2.4 g/L  
NaCl 8.76 g/L  
Adjust pH to 7.4 with HCl
4. 1% BSA-TBS
 

Add 10 g/L of BSA to TBS  
Keep aliquots frozen at -20 °C, or use fresh.
5. Uranyl acetate
 

Prepare a saturated solution of uranyl acetate in 50% ethanol. Filter it through a filter paper. Use fresh.
6. Lead citrate
  - a. Dissolve 1.33 g  $\text{Pb}(\text{NO}_3)_2$  and 1.76 g  $\text{Na}_3(\text{C}_6\text{H}_5\text{O}_7) \cdot 2\text{H}_2\text{O}$  in 30 ml of deionized water. Shake it for 30 min in the orbital shaker. At this stage the solution may appear milky.
  - b. Add 8 ml of 1 M NaOH: the solution should become transparent. Adjust the volume to 50 ml with deionized water. pH should be about 12.  
Keep in small aliquots at -20 °C. Thaw and spin down before use.

### **Acknowledgments**

This protocol was adapted from Sinetova *et al.* (2012). We appreciate Prof. J. Moroney (Louisiana State University, USA) for providing the strain of *C. reinhardtii* 137mt+ and CCM-mutant *cia3*, Prof. G. Samuelsson (Umea University, Sweden) who kindly supplied us with primary antibodies against recombinant Cah3 and Cah3-MBP fusion protein. We thank Prof. Pronina N.A. and Dr. Kupriyanova E.V. for the physiological and molecular biological base of this protocol. This work was supported by a grant from the Russian Foundation for Basic Research (no. 10-04-01463), a grant from the Presidium of Russian Academy of Sciences (program “Molecular and Cell Biology”), and a grant from the Russian Ministry of Education and Science (no. 16.740.11.0176).

### **References**

1. Oliver, C. (2010). [Postembedding labeling methods](#). *Methods Mol Biol* 588: 387-395.

2. Reynolds, E. S. (1963). [The use of lead citrate at high pH as an electron-opaque stain in electron microscopy](#). *J Cell Biol* 17: 208-212.
3. Sinetova, M. A., Kupriyanova, E. V., Markelova, A. G., Allakhverdiev, S. I. and Pronina, N. A. (2012). [Identification and functional role of the carbonic anhydrase Cah3 in thylakoid membranes of pyrenoid of \*Chlamydomonas reinhardtii\*](#). *Biochim Biophys Acta* 1817(8): 1248-1255.