

## MTT Assay of Cell Numbers after Drug/Toxin Treatment

Ran Chen\*

Department of Genetics, Stanford University, Stanford, USA

\*For correspondence: [rcchen@jfkbio.com](mailto:rcchen@jfkbio.com)

**[Abstract]** MTT assay is a colorimetric method for measuring the activity of enzymes in living cells that reduce MTT to formazan dyes, giving a purple color. It is commonly used to determine cytotoxicity of potential medicinal agents and toxic materials, since these types of materials are expected to stimulate or inhibit cell viability and growth. Here, a general protocol is described to carry out an MTT assay on different types of cells.

### Materials and Reagents

1. Raw264.7, MCF-7 or HeLa cells
2. Thiazolyl blue tetrazolium bromide (MTT) (Sigma-Aldrich, catalog number: M5655)
3. DMSO
4. DPBS (Life Technologies, Invitrogen™, catalog number: 14190-250)
5. General chemicals (Sigma-Aldrich)

### Equipment

1. 96 well plate
2. Shaking table
3. Paper towels
4. Incubator

### Procedure

1. Plate 500-10,000 cells in 200  $\mu$ l media per well in a 96 well plate. Leave 8 wells empty for blank controls.
2. Incubate (37 °C, 5% CO<sub>2</sub>) overnight to allow the cells to attach to the wells.
3. Add 2  $\mu$ l of drug of interest dissolved in DMSO to each well. Place on a shaking table, 150 rpm for 5 min, to thoroughly mix the samples into the media.
4. Incubate (37 °C, 5% CO<sub>2</sub>) for 1-5 days to allow the drug/toxin to take effect.

5. Make 2 ml or more of MTT solution per 96 well plate at 5 mg/ml in DPBS. Do not make a stock as MTT in solution is not stable long-term.
6. Add 20  $\mu$ l MTT solution to each well. Place on a shaking table, 150 rpm for 5 min, to thoroughly mix the MTT into the media.
7. Incubate (37 °C, 5% CO<sub>2</sub>) for 1-5 h to allow the MTT to be metabolized.
8. Dump off the media (dry plate on paper towels to remove residue if necessary).
9. Resuspend formazan (MTT metabolic product) in 200  $\mu$ l DMSO. Place on a shaking table, 150 rpm for 5 min, to thoroughly mix the formazan into the solvent.
10. Read optical density at 560 nm and subtract background at 670 nm. Optical density should be directly correlated with cell quantity.

### **Acknowledgments**

This work was funded by 5050 project by Hangzhou Hi-Tech District, Funding for Overseas Returnee by Hangzhou City, ZJ1000 project by Zhejiang Province. This protocol was developed in the Cohen Lab, Department of Genetics, Stanford University, CA, USA [Chen *et al.* (unpublished)].

### **References**

1. Hayon, T., Dvilansky, A., Shpilberg, O. and Nathan, I. (2003). [Appraisal of the MTT-based assay as a useful tool for predicting drug chemosensitivity in leukemia](#). *Leuk Lymphoma* 44(11): 1957-1962.