

Quantification of Cell Corpses, Cell Death Occurrence, Cell Corpse Duration

Yan Zhang, Haibin Wang Xiaochen Wang*

National Institute of Biological Sciences, Zhongguancun Life Science Park, Beijing, China

*For correspondence: wangxiaochen@nibs.ac.cn

[Abstract] During the development of the *C. elegans* hermaphrodite, 131 of the 1090 somatic cells generated undergo programmed cell death, among which 113 die during embryogenesis starting from 200-cell stage. The apoptotic cells (also called “cell corpses”) appear as highly refractile button-like objects and are easily identified using differential interference contrast (DIC) optics (Robertson *et al.*, 1982).

Materials and Reagents

1. Agar pad (made by melting and coating 4% agar on glass slides)
2. *C. elegans* strains [Wild type (N2), Engulfment-defective mutants: commonly used mutants including *ced-1*, *ced-5*, *ced-7*, *ced-6*, *ced-2*, *ced-12*, *ced-10*, which all contain persistent cell corpses that are easily detected under DIC optics]
3. KH_2PO_4
4. Na_2HPO_4
5. NaCl
6. MgSO_4
7. Peptone
8. Cholesterol
9. Vaseline
10. NGM agar (see Recipes)
11. M9 (used to mount embryos on agar pads) (see Recipes)

Equipment

1. Zeiss Axioimager M1 microscope (Zeiss)
2. AxioCam monochrome digital camera (Zeiss)
3. AxiovisionRel 4.7 software (Zeiss)
4. Slides
5. Coverslips

Procedure

A. Quantification of cell corpses

C. elegans is grown on NGM agar plates carrying a lawn of bacterial and kept at 20 °C. Embryos are randomly picked from NGM agar plates containing mix-staged worms and mounted on slides with agar pads in M9 and covered with coverslips. Cell corpses are observed using a Zeiss Axioimager M1 microscope equipped with DIC at 20 °C. Cell corpses are identified by the “raised button” morphology and quantified in the head region of living embryos either at the six different embryonic stages (comma, 1.5-fold, 2-fold, 2.5-fold, 3-fold, 4-fold) for a time course analysis (Figure 2) or at the 4-fold embryonic stage. 15 embryos are counted at each embryonic stage for each strain.




Figure 1. Seven Apoptotic cells (arrows) which appear as “raised button” objects in a *C. elegans* embryo are shown. Some apoptotic cells are not visible at this focus plane.




Figure 2. Time-course analysis of cell corpses during embryonic development was performed in wild-type (N2, open bar), *nrf-5(qx16)* (black bar), or *nrf-5(sa513)* (gray bar). At least 15 embryos were scored at each stage. Data are shown as mean+SEM. Data derived from N2 and *nrf-5(qx16)* or N2 and *nrf-5(sa513)* were compared by unpaired t test. **P < 0.0001; all other points had P > 0.05.

B. Monitor the occurrence of embryonic cell death and cell corpse duration

C. elegans embryos at the two-cell stage are mounted on slides with agar pads in M9 and coverslips are sealed with Vaseline. Images in a 20 micron z series (0.5 micron per section) were captured every 1 min for 8 h using a Zeiss Axioimager M1 microscope equipped with an AxioCam monochrome digital camera. Images are processed and viewed using AxiovisionRel 4.7 software. Embryonic cell deaths are followed during 200-370 min after the first embryonic cleavage by the appearance of the “raised button” morphology of cell corpses. The duration of cell corpses is determined by following the appearance and disappearance (no button-like morphology can be seen) of apoptotic cells. At least three embryos from each strain are followed and quantified. The standard error of the mean (SEM) is used as y error bars for bar charts plotted from the mean value of the data. Data derived from different genetic backgrounds were compared by Student’s two way unpaired t-test. Data were considered statistically different at P < 0.05.

Recipes

1. M9
 - 0.022 M KH₂PO₄
 - 0.042 M Na₂HPO₄
 - 0.086 M NaCl
 - 0.001 M MgSO₄

2. NGM agar
 - NaCl 3 g
 - Agar 17 g
 - Peptone 2.5 g
 - Cholesterol (5 mg/ml in EtOH) 1 ml
 - H₂O 975 ml

Acknowledgments

This protocol is adapted from Robertson *et al.* (1982) and Stanfield and Horvitz (2000).

References

1. Robertson, A. M. G. and Thomson, J. N. (1982). [Morphology of programmed cell death in the ventral nerve cord of *Caenorhabditis elegans* larvae.](#) *J Embryol Exper Morphol* 67(1): 89-100.
2. Stanfield, G. M. and Horvitz, H. R. (2000). [The *ced-8* gene controls the timing of programmed cell deaths in *C. elegans*.](#) *Mol Cell* 5(3): 423-433.