

***Shigella* IpaD and IpaB Surface Localizations**

Lionel Schiavolin*, Alaeddine Meghraoui and Abdelmounnaim Allaoui

Molecular Bacteriology Lab, Université Libre de Bruxelles, Anderlecht, Belgium

*For correspondence: lschiavo@ulb.ac.be

[Abstract] *Shigella* uses a type III secretion system to invade host cell and to cause disease. Secretion control and insertion of a translocation pore into cell membrane are critical steps for pathogenesis and are tightly linked to the formation of the needle tip complex formed by the IpaB and IpaD proteins (Veenendaal *et al.*, 2007). Surface localizations of IpaD and IpaB were monitored by FACS analysis according to the localization protocol for *Pseudomonas aeruginosa* homolog PcrV (Lee *et al.*, 2010).

Materials and Reagents

1. *Shigella* strains
2. Tryptic Soy Broth (TSB) (VWR International, catalog number: for Europe 1.00525.5000 and for U.S. EM1.00525.5007)
3. Agar (MP Biomedicals, catalog number: 0210026291)
4. Congo Red (VWR International, for Europe catalog number: 34140.184)
5. Anti-IpaD and -IpaB polyclonal antibodies (house made)
6. PBS (Fischer Scientific, catalog number: BP399-20)
7. Paraformaldehyde (PFA) (Sigma-Aldrich, catalog number: P6148)
8. Triton X-100 (VWR International, catalog number: 1.08603.1000)
9. Trizma base (Sigma-Aldrich, catalog number: T1503)
10. Bovine serum albumin (BSA) (VWR International, catalog number: 422361V)
11. Anti-mouse secondary antibody CF647-conjugated (Sigma-Aldrich, catalog number: SAB4600351)
12. PBS + PFA 4% stock solution (see Recipes)
13. Congo Red agar plates (see Recipes)

Equipment

1. 250 ml Erlenmeyer
2. Microtubes
3. CR agar plate

4. Polystyrene tube for FACS analysis
5. Centrifuge Heating magnetic stirrer
6. 37 °C shaker
7. Rotator mixer
8. Flow cytometry with a four-colour FACS Calibur cytometer (Becton Dickinson and Company)

Procedure

A. Preparation of bacterial samples

1. Launch overnight precultures in TSB (37 °C with shaking) from Congo Red (CR) positives colonies of *Shigella* on CR agar plates.
2. Dilute 1:100 precultures in fresh TSB (for volume see step A3) and incubate at 37 °C with shaking until an OD₆₀₀ ≈ 1.5 is reached. (Medium must be filtered or autoclaved with stirring to avoid glucose caramelization which interferes with type III secretion.)
3. Harvest 2 x 10⁸ bacteria per tested conditions at 2,000 x g for 4 min at room temperature (RT) (OD₆₀₀ of 1 correspond to approximately 5 x 10⁸ bacteria).
4. Wash twice with 500 µl of ice-cold PBS with 0.1% Triton X-100 (centrifugation conditions as in step A3). Be careful when you take out the liquid supernatant as the pellet can detach gradually all along the procedure.
5. Resuspend in 500 µl of ice-cold PBS with 0.1% Triton X-100 and add 500 µl of PBS + PFA (4%) to fix bacteria.
6. Mix by inversion and incubate 20 min at RT.
7. Add 50 µl Tris HCl 1 M (pH 7.5), mix by inversion and incubate 5 min at RT to quench the cross-linker.
8. Harvest bacteria at 10,000 x g for 2 min at RT (all further centrifugation steps are performed with these parameters).
9. Wash once with 1 ml PBS + 0.1% Triton X-100 and once with 1 ml PBS.

B. Immunological staining

1. Harvest bacteria and resuspend in 500 µl of the blocking solution (PBS + 4% BSA).
2. Incubate bacteria on rotator mixer 1 h at 4 °C.
3. Harvest bacteria and resuspend in 250 µl PBS + 4% BSA with mouse sera diluted 1:100. Sera were from Swiss mice immunized with GST-IpaD¹³¹⁻³³² (Schiavolin *et al.*, 2013) or His-IpgC + IpaB (Page *et al.*, 1999). Negative controls for IpaD and IpaB localizations are respectively an IpaD protein lacking its last residues which do not bind the needle tip (Espina *et al.*, 2006) and an *ipad* KO mutant where IpaB is not retained at the tip

(Veenendaal *et al.*, 2007). In both cases secreted IpaD or IpaB do not interfere with the assay.

4. Incubate overnight at 4 °C with agitation.
5. Add 750 µl PBS to bacteria and then wash twice with 1 ml PBS. All further steps are made in the dark to preserve fluorescent properties of the secondary antibody. Microtubes or rack are covered with aluminum foil.
6. Resuspend in 250 µl of PBS + 4% BSA with goat CF647-conjugated anti-mouse IgG antibody diluted 1:500.
7. Agitate 1 h or longer at 4 °C (incubation time may last overnight).
8. Add 750 µl PBS to bacteria and then wash twice with 1 ml PBS.
9. Resuspend bacterial pellet in 500 µl of PBS and dilute 1:10 in FACS polystyrene tube.
10. Analyze by flow cytometry with a four-colour FACS Calibur cytometer for instance (you will find examples of results in the supporting information of the third reference (Schiavolin *et al.*, 2013) available for free on *Mol. Microbiol.* website).
11. Parameters used (no compensation):

detectors	voltage	mode
FSC	E02	Log
SSC	366	Log
FL4	800	Log

Recipes

1. PBS + PFA 4% stock solution (100 ml)
 - Weigh out 4 g of paraformaldehyde in a 250 ml erlenmeyer
 - Add 80 ml of double-distilled water
 - Add 50 µl of 1 M NaOH
 - Add a magnetic bar and close the erlenmeyer
 - Heat at 70 °C until complete solubilization (the solution should become clear)
 - Put on ice and allow the solution to cool down to RT
 - Adjust volume to 90 ml with double-distilled water
 - Add 10 ml of PBS 10x and mix
 - Filter solution with a 0.2 µm 25 mm nylon syringe filter and aliquot in 15 ml falcon tube
 - Freshly prepared PFA can be stored at -20 °C for further assays (thaw gently at RT)
2. Congo Red agar plates
 - Prepare a 30 g/l TSB solution with bidistilled water and dissolve 15 g/l agar
 - Autoclave the medium for 15 min at 120°C (if you autoclave your medium for 20 min, sugar will caramelize and your plates will be darker, and even darkest after incubation with bacteria at 37°C)

Add the Congo Red at a final concentration of 250 µg/ml (stock solution is prepared in bidistilled water at a concentration of 10 mg/ml) when the bottle can be safely handled with a protective glove.

Acknowledgments

This protocol was adapted from Lee *et al.* (2010). This study was supported by grants from the Belgian Fonds National de la Recherche Scientifique Médicale (FRS-FNRS; Convention 3.4556.11) and from the European Community's Seventh framework program FP7/2011–2015 under grant agreement No. 261742. L.S. and A.M are recipients of a PhD fellowship from the Belgian Fonds National de Recherches Industrielles et Agronomiques (FRIA). A part of this work was also supported by the Fonds Defay, and by Van Buuren and Héger-Masson foundations. We thank A. Op de Beeck for her help in performing the FACS experiment.

References

1. Espina, M., Olive, A. J., Kenjale, R., Moore, D. S., Ausar, S. F., Kaminski, R. W., Oaks, E. V., Middaugh, C. R., Picking, W. D. and Picking, W. L. (2006). [IpaD localizes to the tip of the type III secretion system needle of *Shigella flexneri*](#). *Infect Immun* 74(8): 4391-4400.
2. Lee, P. C., Stopford, C. M., Svenson, A. G. and Rietsch, A. (2010). [Control of effector export by the *Pseudomonas aeruginosa* type III secretion proteins PcrG and PcrV](#). *Mol Microbiol* 75(4): 924-941.
3. Page, A. L., Ohayon, H., Sansonetti, P. J. and Parsot, C. (1999). [The secreted IpaB and IpaC invasins and their cytoplasmic chaperone IpgC are required for intercellular dissemination of *Shigella flexneri*](#). *Cell Microbiol* 1(2): 183-193.
4. Schiavolin, L., Meghraoui, A., Cherradi, Y., Biskri, L., Botteaux, A. and Allaoui, A. (2013). [Functional insights into the *Shigella* type III needle tip IpaD in secretion control and cell contact](#). *Mol Microbiol* 88(2): 268-282.
5. Veenendaal, A. K., Hodgkinson, J. L., Schwarzer, L., Stabat, D., Zenk, S. F. and Blocker, A. J. (2007). [The type III secretion system needle tip complex mediates host cell sensing and translocon insertion](#). *Mol Microbiol* 63(6): 1719-1730.